


# Hypertensives may benefit from gut microbial metabolites, study finds

January 10 2023



HAMSAB promotes the expansion of SCFA-producing commensal bacteria. 16S rRNA sequences were obtained from fecal samples and analyzed using QIIME2. a, Chao1, observed and Shannon  $\alpha$ -diversity indices were analyzed using

MicrobiomeAnalyst software comparing the first and second baselines (n = 40 samples, two-tailed paired t-test). b, Principal coordinate analysis (PCoA) plot showing weighted Bray–Curtis index at baseline of participants randomized to HAMSAB or placebo diet (n = 40 samples). c, Chao1, observed and Shannon  $\alpha$ -diversity indices comparing HAMSAB and placebo diet (n = 39 samples, two-tailed paired t-test). d, PCoA plot showing  $\beta$ -diversity measured by Bray–Curtis index of baselines of participants after HAMSAB or placebo diet (n = 39 samples). Filtered count and log-transformed count showing relative abundances of *P. distasonis* (e) and *R. gausvrauii* (f) (n = 39 samples) (edgeR analysis). Box plots in a, c, e and f show the median and interquartile range of each group. Shaded ellipsis in PCoA plots represent the 95% CI for each group. Credit: *Nature Cardiovascular Research* (2023). DOI: 10.1038/s44161-022-00197-4

Hypertension, or high blood pressure, is a leading cause of death worldwide.

Four out of five hypertensive patients have uncontrolled [blood pressure](#) despite clinical and lifestyle therapy, suggesting new approaches are needed.

Over the past decade, [experimental research](#) has shown that gut microbiota and the substances they produce regulate blood pressure.

Dietary fiber, especially fermentable fiber, is digested by commensal gut microorganisms in the [large intestine](#), modulating the gut microbiota and releasing substances called short-chain [fatty acids](#) (SCFAs), such as acetate and butyrate.

SCFAs regulate host physiology through bioactive pathways. In animal models, SCFA administration decreases blood pressure.

Previous studies led by Associate Professor Francine Marques from the

Monash University School of Biological Sciences had found that acetate and butyrate supplementation [lower blood pressure](#) in hypertension models.

A new randomized clinical trial led by Associate Professor Marques, published Jan. 9 in *Nature Cardiovascular Research*, shows that SCFAs can also lower blood pressure of hypertensive patients.

"These metabolites have untapped translational potential," said Associate Professor Marques.

"Our previous studies found that acetate and butyrate, two microbial SCFAs, decreased blood pressure in mice. However, this method would require patients to ingest SCFAs 24/7, making it unsuitable for humans."

Thus, in the clinical trial, her research team used acetylated and butyrylated high amylose maize (HAMSAB), a fermentable fiber that produces large quantities of acetate and butyrate.

They recruited 20 untreated hypertensives for a 3-week double-blind, placebo-controlled, randomized cross-over phase II trial of HAMSAB to lower blood pressure.

The team assessed home and 24-hour blood pressure, plasma SCFAs, fecal microbiota, gastrointestinal transit, pH, and cytokines.

"In HAMSAB-treated hypertensives, 24-hour [systolic blood pressure](#) dropped 6.1 mmHg," Associate Professor Marques said. "This is equivalent to one blood pressure lowering drug and has important clinical implications."

Further analysis showed that plasma acetate and butyrate were considerably higher after the three-week HAMSAB intervention than the

placebo arm. "The levels of SCFAs we were able to achieve with HAMSAB were remarkable and higher than can be achieved with diet alone," said Dr. Hamdi Jama, the lead author of the study.

HAMSAB also altered gut microbial makeup, favoring acetate- and butyrate-producing bacteria like *Ruminococcus* species.

"Our clinical trial shows acetate and butyrate, gut microbial metabolites, reduce blood pressure in patients, suggesting medicinal potential," Dr. Jama said.

"This supports clinical and experimental findings that [gut microbiota](#) alterations, particularly SCFA-producer depletion, may predate hypertension," Associate Professor Marques said.

"Thus, fermentable fibers like HAMSAB may re-establish gut microbial communities that produce SCFAs. This approach may aid in the reduction of rates of hypertension and cardiovascular disease worldwide."

Associate Professor Marques said HAMSAB lowers blood pressure, but multi-center trials with larger samples and longer follow-up could provide more conclusive results.

**More information:** Hamdi A. Jama et al, Prebiotic intervention with HAMSAB in untreated essential hypertensive patients assessed in a phase II randomized trial, *Nature Cardiovascular Research* (2023). [DOI: 10.1038/s44161-022-00197-4](https://doi.org/10.1038/s44161-022-00197-4)

Provided by Monash University

Citation: Hypertensives may benefit from gut microbial metabolites, study finds (2023, January 10) retrieved 28 December 2023 from <https://medicalxpress.com/news/2023-01-hypertensives-benefit-gut-microbial-metabolites.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.