

Nitrite additives associated with increased risk of type 2 diabetes

January 17 2023



Credit: Pixabay/CC0 Public Domain

Nitrites and nitrates occur naturally in water and soil and are commonly ingested from drinking water and dietary sources. They are also used as food additives to increase shelf life. A study publishing January 17 in the

open access journal *PLOS Medicine* by Bernard Srour of the Nutritional Epidemiology Research Team (EREN-CRESS) of Inserm, INRAE, Cnam, and Sorbonne Paris Nord University, Bobigny, France, and colleagues suggests an association between dietary exposure to nitrites and risk of type 2 diabetes.

Some public health authorities have advocated for limiting the use of nitrites and [nitrates](#) as food additives. However, the role of dietary nitrites and nitrates in metabolic dysfunction and type 2 diabetes in humans remains unexplored. In order to investigate the relationship between dietary exposure to nitrites/nitrates type 2 diabetes risk, researchers accessed data collected from 104,168 participants in the prospective cohort NutriNet-Santé.

The NutriNet-Santé study is an ongoing, web-based cohort study initiated in 2009. Participants aged 15 and older enroll voluntarily and self-report [medical history](#), sociodemographic, diet, lifestyle, and major health updates. The researchers used detailed nitrite/nitrate exposure, derived from several databases and sources, and then developed statistical models to analyze self-reported diet information with [health outcomes](#).

The researchers found that participants in the NutriNet-Santé cohort reporting a higher intake of nitrites overall and specifically from food additives, and non-additives sources had a higher risk of developing type 2 diabetes. There was no association between nitrates and type 2 diabetes risk, and the findings did not support any [potential benefits](#) for dietary nitrites or nitrates in terms of protection against type 2 diabetes.

The study had several limitations and additional research is required to validate the results. The data were self-reported and the researchers could not confirm specific nitrite/nitrate exposure using biomarkers due to the underlying biological challenges. Additionally, people in the

cohort's demographics and behaviors may not be generalizable to the rest of the population—the cohort included a greater number of younger individuals, more often women, who exhibited healthier behaviors. Residual confounding may also have impacted the outcomes as a result of the observational design of the study.

According to the authors, "These results provide a new piece of evidence in the context of current discussions regarding the need for a reduction of nitrite additives' use in processed meats by the food industry, and could support the need for better regulation of soil contamination by fertilizers. In the meantime, several public health authorities worldwide already recommend citizens to limit their consumption of foods containing controversial additives, including sodium [nitrite](#)."

Srouf and Touvier add, "This is the first largescale cohort study to suggest a direct association between additives-originated nitrites and type-2 [diabetes](#) risk. It also corroborates previously suggested associations between total dietary nitrites and T2D risk."

More information: Dietary exposure to nitrites and nitrates in association with type 2 diabetes risk: Results from the NutriNet-Santé population-based cohort study, *PLoS Medicine* (2023). [DOI: 10.1371/journal.pmed.1004149](#)

Provided by Public Library of Science

Citation: Nitrite additives associated with increased risk of type 2 diabetes (2023, January 17) retrieved 27 March 2023 from <https://medicalxpress.com/news/2023-01-nitrite-additives-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.