


Is the brain wired differently in people with addiction?

February 16 2023, by Mylène Tremblay



Credit: Pixabay/CC0 Public Domain

Some 10% to 15% of people will have a substance abuse problem at some point in their life, making it one of the most common psychiatric disorders. Risks associated with substance abuse include dangerous driving, missed work, problems with depression, anxiety, health and money problems. Yet people with addictions seem to experience a sort of myopia, getting in deeper and deeper to cope with unpleasant emotions and to feel good, regardless of the cost.

Research has clearly established that psychosocial and [environmental factors](#) play a role in the development of [substance abuse](#) disorders. But there is more to the story, and now [functional magnetic resonance](#) imaging (MRI) studies are highlighting the importance of biological factors, in particular how the brain functions.

"The influence of such factors can be as much as 50%—this astonishingly high figure is why we're so interested in what's going on in the brains of substance abusers," explained Stéphane Potvin, a professor in Université de Montréal's Department of Psychiatry and Addiction whose research focuses primarily on the harmful effects of cannabis and alcohol on the brain structures of people with schizophrenia.

Zeroing in on markers

Potvin, who coordinates the Neurobiology and Mental Health division of the Centre de recherche de l'Institut universitaire en santé mentale de Montréal, wanted to zero in on the neurobiological markers associated with addiction. With his student Jules R. Dugré, a Ph.D. candidate in [biomedical sciences](#), he conducted a [meta-analysis](#) of 96 studies involving a total of 5,757 subjects with some form of substance dependence—alcohol, nicotine, cannabis, psychostimulants or another drug.

Rather than focusing on [brain activity](#) as a whole, in his analysis Potvin

and Dugré looked at what is known as "functional connectivity," in other words, how well different parts of the brain communicate with each other. This is the first time an analysis of this type has been done.

Published in November in *Addiction Biology*, the study reveals a number of anomalies in the brain's reward, decision-making and habit formation systems:

1. Reward

"The brains of substance abusers tend to show hyperconnectivity between the [ventromedial prefrontal cortex](#) and [ventral striatum](#), two key areas of the reward system," said Potvin. "This could explain the tendency to choose immediate gratification and why the substance's motivational value increases over time despite the negative impact on other spheres of life."

2. Decision-making

"We also see reduced connectivity in certain [brain areas](#) involved in decision-making, such as the prefrontal cortex and amygdala," added Potvin. "This is consistent with the fact that substance abusers can seem indifferent to the harmful consequences of their choices."

3. Habit formation

The most original finding is the presence of anomalies in the brain regions associated with habit formation, including the dorsal striatum and premotor cortex. "This hyperconnectivity could explain the compulsive nature of substance abuse," Potvin noted.

No charges in impulse control

The meta-analysis did not, however, show changes in the [brain regions](#) and networks associated with impulse control. This contrasts with previous studies, which showed abnormalities in these regions using other neuroimaging approaches.

Potvin is quick to point out the limitations of his study. "In our analysis, we included all relevant studies of substance abusers, whatever the substance. But are brain systems equally affected by different substances? We really don't know."

Ultimately, these findings will help guide the development of neuromodulation-based interventions in the field of addiction.

"First we need a better understanding of the brain systems involved in substance abuse," said Potvin. "This will allow us to pinpoint the areas to stimulate or inhibit in order to rebalance the brain and change behavior. The more solid evidence we have, the more treatment centers will be motivated to invest in the equipment needed to carry out this type of intervention."

More information: Jules R. Dugré et al, Disrupted functional connectivity of the brain reward system in substance use problems: A meta-analysis of functional neuroimaging studies, *Addiction Biology* (2022). [DOI: 10.1111/adb.13257](https://doi.org/10.1111/adb.13257)

Provided by University of Montreal

Citation: Is the brain wired differently in people with addiction? (2023, February 16) retrieved 14 February 2024 from <https://medicalxpress.com/news/2023-02-brain-wired-differently-people-addiction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.