


Genetic factors contributing to connection issues for white matter in the human brain discovered

February 20 2023, by Bob Yirka



SNP-based heritability and mvGWAS analyses of node-level connectivity and edge-level connectivity in 30,810 participants.(**A**) All 90 node-level (i.e., regional) connectivities showed significant SNP-based heritability after Bonferroni correction, ranging from 7.8 to 29.5%. (**B**) Eight hundred fifty-one of 947 edge-level connectivities showed significant SNP-based heritability after Bonferroni correction, ranging from 4.6 to 29.5%. Right: Brain maps. Left:



Nodes grouped by frontal, prefrontal, parietal, temporal, and occipital cortical lobes and subcortical structures. Heritabilities can be visualized interactively in a dynamic Web-based interface (see "Data and materials availability" statement). (C) Miami plot for mvGWAS of 90 node-level connectivities (top) and 851 edge-level connectivities (bottom). The black lines indicate the genome-wide significance threshold *P*

Citation: Genetic factors contributing to connection issues for white matter in the human brain discovered (2023, February 20) retrieved 23 March 2023 from <u>https://medicalxpress.com/news/2023-02-genetic-factors-contributing-issues-white.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.