


Novel technique distinguishes between types of oral bone lesion based on MRI scan image texture

February 15 2023, by Julia Moióli



(A) Ameloblastoma located in the mandibular body on the left side. Arrows represent regions with variable signal intensity, which are characterized by hypersignal representing higher aqueous content, a characteristic of T2-weighted sequences. (B) Segmentation process represented by the red contour. Only the most homogeneous portion of the lesion was segmented, that is, the region shown by the hypersignal. Credit: *Scientific Reports* (2022). DOI: 10.1038/s41598-022-20802-7

Ameloblastomas and odontogenic keratocysts are benign lesions of the maxillo-mandibular region with different biological characteristics. The former are often aggressive, forming a large tumor and growing into the jawbone.

The latter are cysts that can be managed and treated in various ways. However, they have practically identical morphological characteristics and are extremely difficult to distinguish using conventional imaging techniques such as X-rays, MRI scans and CAT scans, making treatment hard to plan and potentially requiring additional procedures.

In search of a solution to predict the type of lesion and facilitate surgical planning, researchers at Cruzeiro do Sul University (UNICSUL), the University of São Paulo (USP) and the State University of Campinas (UNICAMP) in Brazil, in partnership with colleagues at the University of Gothenburg in Sweden and the University of Ankara in Turkey, used an image processing technique called texture analysis for the first time on these lesions. The study is described in an article published in the journal *Scientific Reports*.

"In three-dimensional imaging techniques, such as MRI and CAT scans, the voxels [units representative of a given volume of 3D space] and pixels are arranged differently, with distances and shades of gray that vary depending on the tissue being scanned. This information can be converted into numerical values [algorithms] to produce a purely mathematical and [statistical analysis](#)," said André Luiz Ferreira Costa, last author of the article and a professor in UNICSUL's program of graduate studies in dentistry.

The study sample comprised 18 patients diagnosed with one of the lesions and undergoing treatment at the dentistry division of Hospital das Clínicas, the hospital complex run by USP's Medical School. The diagnosis was confirmed by biopsy in all cases. Eight had

ameloblastomas and ten had keratocysts. The analysis used their MRI scans, which were chosen because they are three-dimensional and do not involve radiation, so that they can be repeated.

Eleven textural parameters were measured at five different distances, for a total of 55 variables, which were processed using MATLAB scientific software. The variables "entropy" and "sum average" were found to be statistically significant. The former refers to the degree of disorder among pixels in the image of interest, while the latter refers to the average of the sums of two-pixel values in the image. Keratocysts display greater uniformity and less gray-level disorder than ameloblastomas.

"The main achievement of the study is the possibility of obtaining a definitive result more quickly by means of imaging and therefore of more suitable and safer treatment," Costa said. "These two types of lesion are extracted differently. Keratocysts are removed and the remaining cavity is thoroughly cleaned. This is a relatively simple surgical procedure. Ameloblastomas require a larger safety margin, involving bone resection because of their aggressive behavior."

Next steps

The results of the study are promising in so far as they show that MRI scan parameter measurement can be used to distinguish between ameloblastomas and keratocysts as an important diagnostic aid. Next steps will include studies involving other types of lesions, such as unicystic ameloblastoma, which is relatively rare, and orthokeratinized odontogenic keratocyst.

The researchers also plan to increase the size of the study sample. Because texture analysis is based on the signal intensity of pairs of pixels, the results will be better if there are more patients in the study

sample and the lesions are larger.

More information: João Pedro Perez Gomes et al, Magnetic resonance imaging texture analysis to differentiate ameloblastoma from odontogenic keratocyst, *Scientific Reports* (2022). [DOI: 10.1038/s41598-022-20802-7](https://doi.org/10.1038/s41598-022-20802-7)

Provided by FAPESP

Citation: Novel technique distinguishes between types of oral bone lesion based on MRI scan image texture (2023, February 15) retrieved 3 July 2023 from <https://medicalxpress.com/news/2023-02-technique-distinguishes-oral-bone-lesion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.