


AI accurately identifies normal and abnormal chest X-rays

March 7 2023



A 44-year-old man who presented with chest pain and dyspnea. (A) Chest X-ray shows very subtle nodular opacities, primarily in lower lobes, representative of pneumonia and a discrete silhouette sign of the right cardiac border (arrow). The AI system interpreted this chest X-ray as normal. It was also interpreted as normal in the clinical radiology report. (B) CT scan shows the lower lobe airspace opacities with vague tree-in-bud morphology (box) and an area of consolidation (arrow). Pulmonary angiography was performed 5 hours after X-ray. This was the sole false-negative “critical” finding by the AI tool. Credit: Radiological Society of North America

An artificial intelligence (AI) tool can accurately identify normal and abnormal chest X-rays in a clinical setting, according to a study published in *Radiology*.


Chest X-rays are used to diagnose a wide variety of conditions to do with the heart and lungs. An abnormal chest X-ray can be an indication of a range of conditions, including cancer and chronic lung diseases.

An AI tool that can accurately differentiate between normal and abnormal chest X-rays would greatly alleviate the heavy workload experienced by [radiologists](#) globally.

"There is an exponentially growing demand for [medical imaging](#), especially cross-sectional such as CT and MRI," said study co-author Louis Lind Plesner, M.D., from the Department of Radiology at the Herlev and Gentofte Hospital in Copenhagen, Denmark. "Meanwhile, there is a global shortage of trained radiologists. Artificial intelligence has shown great promise but should always be thoroughly tested before any implementation."

For this retrospective, multi-center study, Dr. Plesner and colleagues wanted to determine the reliability of using an AI tool that can identify normal and abnormal chest X-rays.

Researchers used a commercially available AI tool to analyze the chest X-rays of 1,529 patients from four hospitals in the capital region of Denmark. Chest X-rays were included from emergency department patients, in-hospital patients and outpatients. The X-rays were classified by the AI tool as either "high-confidence normal" or "not high-confidence normal" as in normal and abnormal, respectively.



Four examples (out of a total of nine) of X-rays classified as “abnormal, unremarkable” by the reference standard (provided independently by two board-certified thoracic radiologists, with a third radiologist in cases of disagreement) but as normal by the AI tool. All X-rays show very subtle and unremarkable findings; these X-rays were all classified as normal by the clinical radiologic report as well. (A) X-ray in a 58-year-old woman with very discrete linear atelectasis in the lingula segment of the left upper lobe (arrow). (B) X-ray in a 61-year-old woman shows presence of a cervical rib on the right side (arrow). (C, D) Images in a 48-year-old woman (C) and 64-year-old woman (D) show

very subtle degeneration in the spine with osteophytes in lower thoracic segments (arrow). Credit: Radiological Society of North America

Two board-certified thoracic (chest) radiologists were used as the reference standard. A third [radiologist](#) was used in cases of disagreements, and all three physicians were blinded to the AI results.

Of the 429 chest X-rays that were classified as normal, 120, or 28%, were also classified by the AI tool as normal. These X-rays, or 7.8 % of all the X-rays, could be potentially safely automated by an AI tool. The AI tool identified abnormal chest X-rays with a 99.1% of sensitivity.

"The most surprising finding was just how sensitive this AI tool was for all kinds of chest disease," Dr. Plesner said. "In fact, we could not find a single chest X-ray in our database where the algorithm made a major mistake. Furthermore, the AI tool had a sensitivity overall better than the clinical board-certified radiologists."

According to the researchers, further studies could be directed toward larger prospective implementation of the AI tool where the autonomously reported chest X-rays are still reviewed by radiologists.

The AI tool performed especially well at identifying normal X-rays of the outpatient group at a rate of 11.6%. This suggests that the AI model would perform especially well in outpatient settings with a high prevalence of normal chest X-rays.

"Chest X-rays are one of the most common imaging examination performed worldwide," Dr. Plesner said. "Even a small percentage of automatization can lead to saved time for radiologists, which they can prioritize on more complex matters."

More information: Louis L. Plesner et al, Autonomous Chest Radiograph Reporting Using AI: Estimation of Clinical Impact, *Radiology* (2023). [DOI: 10.1148/radiol.222268](https://doi.org/10.1148/radiol.222268)

Provided by Radiological Society of North America

Citation: AI accurately identifies normal and abnormal chest X-rays (2023, March 7) retrieved 29 April 2023 from

<https://medicalxpress.com/news/2023-03-ai-accurately-abnormal-chest-x-rays.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.