


Not getting enough sleep could blunt antibody response to vaccination, leaving you more vulnerable to infection

March 13 2023



Insufficient sleep around the day of vaccination is associated with a lower antibody response

A cartoon depicting the effects of insufficient sleep on vaccination. Credit: Spiegel et al.

In reviewing data from previous studies, a team lead by researchers at the University of Chicago and the French National Institute of Health and Medical Research (Inserm) found that individuals who had fewer

than six hours of sleep per night in the days surrounding vaccination had a blunted antibody response. That indicates efforts to promote healthy sleep duration ahead of an immunization could be an easy way to improve vaccine effectiveness. The study was published March 13 in *Current Biology*.

The latest work builds on a [2002 study](#) by members of the team showing that restricting sleep in participants diminished their antibody response to [influenza vaccination](#), leading to about half of the antibody levels seen in controls at 10 days after an inoculation. Their interest in the work was revived during the COVID-19 pandemic lockdowns in 2020, when they began to connect with others who had studied this question and started to pull together the meta-analysis.

Across seven studies, which examined the impact of sleep duration on vaccination against viral illnesses such as influenza and hepatitis, the researchers found that insufficient sleep (defined as under six [hours of sleep](#) per night) in the days surrounding vaccination resulted in a decreased antibody response.

"Insufficient sleep is a behavioral factor that can be corrected before vaccination and may not only strengthen, but also extend, the [vaccine response](#)," said Eve Van Cauter, Ph.D., Professor Emeritus of Medicine at UChicago and senior author on the meta-analysis. "We know that people respond differently to vaccination according to their age, sex, existing medical conditions and other factors that cannot be readily changed. Having an easily modifiable behavior that you can adjust around the time of your appointment gives you something you can control that is likely to improve your body's response."

Importantly, the association was seen only in studies that objectively assessed sleep duration using wearable activity trackers or sleep studies in the laboratory. Self-reported sleep duration was not a predictor of

[vaccine](#) response. The researchers noted that while the association was strong for men, it was weaker and not statistically significant for women. They argue this was likely due to the fact that none of the studies in women accounted for variations in sex hormone levels by menstrual cycle, use of contraceptives and menopausal status.

"The link between sleep and [vaccine effectiveness](#) could be a major concern for people with irregular work schedules, especially for shift workers who typically have reduced sleep duration," said Van Cauter. "This is something people should consider planning around, to ensure that they are getting enough sleep in the week before and after their vaccines."

Using the results of the [meta-analysis](#) and comparing to known data on the antibody response to the Pfizer-BioNTech COVID-19 vaccine, the researchers estimated that the effects of insufficient sleep on the vaccine response would be equivalent to two months of waning antibodies after vaccination.

"Interestingly, we saw the same outcomes in both influenza, which is a respiratory virus, and hepatitis, which affects the liver, suggesting that this effect could extend to all kinds of viruses, including coronaviruses like SARS-CoV-2," said Karine Spiegel, Ph.D., first author on the study and a research scientist at Inserm. "Overall, we see these results as a call to action."

The researchers hope that the study will inspire more research into the phenomenon to clarify the effects on men and women, as well as to better understand how different vaccines may be impacted by sleep duration and how sleep might be optimized to promote a better vaccine response.

"We need much larger studies that control for the sex hormone

environment in women in particular," said Spiegel. "We also need a better definition of how many days of short sleep duration affect the antibody response, and whether it is just before the vaccine, or also during and after. Large scale studies that consider behavioral, demographic and hormonal characteristics should provide new insights that will translate to measurable impacts on vaccine efficacy."

"The [immune system](#) is not the only one modulated by sleep," said Van Cauter. "Insufficient sleep is linked to other [health issues](#) such as an increased risk of developing obesity, diabetes or hypertension. Vaccines are an important tool for preventing and reducing the impacts of infectious diseases, and we think that you may be able to implement a simple behavioral change—getting enough sleep—to derive an immediate benefit. It's cheap, and there is no adverse effect."

More information: Karine Spiegel, A meta-analysis of the associations between insufficient sleep duration and antibody response to vaccination, *Current Biology* (2023). [DOI: 10.1016/j.cub.2023.02.017](https://doi.org/10.1016/j.cub.2023.02.017). [www.cell.com/current-biology/f ... 0960-9822\(23\)00156-2](https://www.cell.com/current-biology/fulltext/S0960-9822(23)00156-2)

Provided by University of Chicago Medical Center

Citation: Not getting enough sleep could blunt antibody response to vaccination, leaving you more vulnerable to infection (2023, March 13) retrieved 22 February 2024 from <https://medicalxpress.com/news/2023-03-blunt-antibody-response-vaccination-vulnerable.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--