


# Study: COVID-19 mRNA booster vaccinations increased and sustained antibody responses

March 23 2023



Data at 7 months after the second vaccination (n = 1,138), and at 4 months (n = 701) and 7 months (n = 377) after the third vaccination were used for analysis. The boxplots show medians (middle line) and first and third quartiles (boxes),

while the whiskers show  $1.5 \times$  the interquartile range (IQR) above and below the box. Circles indicate outlier values. Broken lines indicate the cutoff value for the antibody level (10 arbitrary units [AU]/mL). Credit: TMIMS

Booster vaccinations against SARS-CoV-2 are being promoted worldwide to counter the coronavirus disease 2019 (COVID-19) pandemic. Recent studies on the effectiveness of the third dose of the vaccine found improved efficacy against SARS-CoV-2 infection and a reduction in hospitalization. However, most studies on the effect of the third vaccination have been limited to a relatively short period of time after the third dose, and few longitudinal analyses of the effects of the third vaccination with an anti-SARS-CoV-2 vaccine have been reported.

Researchers from Tokyo Metropolitan Institute of Medical Science analyzed the longitudinal effect of the third SARS-CoV-2 mRNA vaccination on antibody responses in [healthcare workers](#) in Tokyo, Japan. Additionally, antibody responses induced by the fourth vaccination were analyzed.

The levels of anti-spike (S) IgG and neutralizing antibody against SARS-CoV-2 were measured at 7 months after the second vaccination ( $n = 1,138$ ), and at 4 ( $n = 701$ ) and 7 ( $n = 417$ ) months after the third vaccination using an iFlash 3000 chemiluminescence immunoassay analyzer (YHLO).

Among the 417 participants surveyed at 7 months after the third vaccination, 40 had received the fourth vaccination. A multiple linear regression analysis was performed to clarify which factors were associated with the anti-S IgG and neutralizing antibody. Variables assessed included sex, age, number of days after the second or third vaccination, diagnostic history of COVID-19, and anti-nucleocapsid (N)

IgG level.

At 7 months after the third vaccination, antibody responses were significantly higher than those at the same time after the second vaccination. Unlike the second vaccination, age had no effect on the antibody responses induced by the third vaccination. Furthermore, the fourth vaccination resulted in a further increase in antibody responses. The multiple linear regression analysis identified anti-N IgG level, presumably associated with infection, as a factor associated with antibody responses.

This work demonstrate that strong and persistent [antibody responses](#) were induced after SARS-CoV-2 mRNA booster vaccinations. These data are expected to be useful for vaccination scheduling.

The paper is published in the *Journal of Infection and Chemotherapy*.

**More information:** Takahiro Sanada et al, Antibody response to third and fourth BNT162b2 mRNA booster vaccinations in healthcare workers in Tokyo, Japan, *Journal of Infection and Chemotherapy* (2022). [DOI: 10.1016/j.jiac.2022.12.012](https://doi.org/10.1016/j.jiac.2022.12.012)

Provided by Tokyo Metropolitan Institute of Medical Science

Citation: Study: COVID-19 mRNA booster vaccinations increased and sustained antibody responses (2023, March 23) retrieved 13 February 2024 from <https://medicalxpress.com/news/2023-03-covid-mrna-booster-vaccinations-sustained.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.