


New study reveals novel insights on brain development sequence through adolescence

April 10 2023



Credit: CC0 Public Domain

Brain development does not occur uniformly across the brain, but follows a newly identified developmental sequence, according to a new Penn Medicine study. Brain regions that support cognitive, social, and emotional functions appear to remain malleable—or capable of changing, adapting, and remodeling—longer than other brain regions, rendering youth sensitive to socioeconomic environments through adolescence. The findings were published recently in *Nature*

Neuroscience.

Using magnetic resonance imaging (MRI), researchers charted how developmental processes unfold across the human brain from the ages of 8 to 23 years old. The findings indicate a new approach to understanding the order in which individual brain regions show reductions in plasticity during development.

Brain plasticity refers to the capacity for [neural circuits](#)—connections and pathways in the brain for thought, emotion, and movement—to change or reorganize in response to internal biological signals or the [external environment](#). While it is generally understood that children have higher brain plasticity than adults, this study provides new insights into where and when reductions in plasticity occur in the brain throughout childhood and adolescence.

The findings reveal that reductions in brain plasticity occur earliest in "sensory-motor" regions, such as visual and auditory regions, and occur later in "associative" regions, such as those involved in higher-order thinking (problem solving and [social learning](#)). As a result, brain regions that support executive, social, and emotional functions appear to be particularly malleable and responsive to the environment during [early adolescence](#), as plasticity occurs later in development.

"Studying [brain development](#) in the living human brain is challenging. A lot of neuroscientists' understanding about brain plasticity during development actually comes from studies conducted with rodents. But rodent brains do not have many of what we refer to as the association regions of the [human brain](#), so we know less about how these important areas develop," said corresponding author Theodore D. Satterthwaite, MD, the McLure Associate Professor of Psychiatry in the Perelman School of Medicine at the University of Pennsylvania, and director of the Penn Lifespan Informatics and Neuroimaging Center (PennLINC).

To address this challenge, the researchers focused on comparing insights from previous rodent studies to youth MRI imaging insights. Prior research examining how neural circuits behave when they are plastic uncovered that brain plasticity is linked to a unique pattern of "intrinsic" brain activity.

Intrinsic activity is the neural activity occurring in a part of the brain when it is at rest, or not being engaged by external stimuli or a mental task. When a brain region is less developed and more plastic, there tends to be more intrinsic activity within the region, and that activity also tends to be more synchronized. This is because more neurons in the region are active, and they tend to be active at the same time. As a result, measurements of brain activity waves show an increase in amplitude (or height).

"Imagine that individual neurons within a region of the brain are like instruments in an orchestra. As more instruments begin to play together in synchrony, the sound level of the orchestra increases, and the amplitude of the sound wave gets higher," said first author Valerie Sydnor, a Neuroscience Ph.D. student.

"Just like decibel meters can measure the amplitude of a sound wave, the amplitude of intrinsic brain activity can be measured with functional MRI while kids are simply resting in the scanner. This allowed our team to study a functional marker of brain plasticity safely and non-invasively in youth."

Analyzing MRI scans from more than 1,000 individuals, the authors found that the functional marker of [brain plasticity](#) declined in earlier childhood in sensory-motor regions but did not decline until mid-adolescence in associative regions.

"These slow-developing associative regions are also those that are vital

for children's cognitive attainment, social interactions, and emotional well-being," Satterthwaite added. "We are really starting to understand the uniqueness of human's prolonged developmental program."

"If a brain region remains malleable for longer, it may also remain sensitive to [environmental influences](#) for a longer window of development," Sydnor said. "This study found evidence for just that."

The authors studied relationships between youths' socioeconomic environments and the same functional marker of plasticity. They found that the effects of the environment on the brain were not uniform across regions nor static across development. Rather, the effects of the environment on the brain changed as the identified developmental sequence progressed.

Critically, youths' socioeconomic environments generally had a larger impact on brain development in the late-maturing associative [brain regions](#), and the impact was found to be largest in adolescence.

"This work lays the foundation for understanding how the environment shapes neurodevelopmental trajectories even through the teenage years," said Bart Larsen, Ph.D., a PennLINC postdoctoral researcher and co-author.

Sydnor elaborated, "The hope is that studying developmental plasticity will help us to understand when environmental enrichment programs will have a beneficial impact on each child's neurodevelopmental trajectory. Our findings support that programs designed to alleviate disparities in youths' socioeconomic environments remain important for brain development throughout the adolescent period."

More information: Valerie J. Sydnor et al, Intrinsic activity development unfolds along a sensorimotor–association cortical axis in

youth, *Nature Neuroscience* (2023). [DOI: 10.1038/s41593-023-01282-y](https://doi.org/10.1038/s41593-023-01282-y)

Provided by Perelman School of Medicine at the University of
Pennsylvania

Citation: New study reveals novel insights on brain development sequence through adolescence (2023, April 10) retrieved 3 February 2024 from <https://medicalxpress.com/news/2023-04-reveals-insights-brain-sequence-adolescence.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.