

Heroin addicts have higher pain sensitivity, even during treatment

April 25 2012



Those taking methadone continue to have heightened pain responses, study finds.

(HealthDay) -- Heroin addicts often have an increased sensitivity to pain, and this sensitivity does not subside over the course of treatment with methadone or other opioids, new research finds.

Researchers from the University of California-Los Angeles sought to determine how increased sensitivity to pain (also called hyperalgesia) might change as a heroin addict moves from drug abuse to stabilization and eventually to maintenance on a pain-treatment opioid such as methadone or buprenorphine.

The investigators assessed the pain responses of 82 heroin addicts who sought treatment and were given either methadone or buprenorphine. The addicts' pain responses were compared to people who didn't use



drugs.

Pain responses in both groups were measured using electrical stimulation and the cold pressor test, in which a hand is placed in ice cold water.

The results showed no significant changes in <u>pain response</u> among heroin addicts who took either methadone or <u>buprenorphine</u> for <u>maintenance</u> <u>therapy</u>.

The study appears in *The Journal of Pain*.

Doctors need to be aware of opioid-induced <u>pain sensitivity</u> associated with addiction in patients prescribed opioids as well as those taking the drugs illegally, according to the study.

More information: The U.S. National Institute on Drug Abuse has more about <u>treatments for drug addiction</u>.

Copyright © 2012 HealthDay. All rights reserved.

Citation: Heroin addicts have higher pain sensitivity, even during treatment (2012, April 25) retrieved 14 February 2024 from https://medicalxpress.com/news/2012-04-heroin-addicts-higher-pain-sensitivity.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.