

Reasons for discontinuation vary by psoriasis treatment

August 10 2012



The reasons for discontinuation of commonly used treatments for psoriasis vary by treatment, according to a study published online July 30 in the *Journal of the American Academy of Dermatology*.

(HealthDay) -- The reasons for discontinuation of commonly used treatments for psoriasis vary by treatment, according to a study published online July 30 in the *Journal of the American Academy of Dermatology*.

Howa Yeung, from the University of Pennsylvania Perelman School of Medicine in Philadelphia, and colleagues conducted interviews of 1,095 [patients](#) with moderate-to-severe plaque psoriasis to assess 11 reasons for treatment discontinuation for all past treatments (2,231 total).

The authors found that the median treatment duration ranged from 6.0 to 20.5 months (P

"A broad range of clinically relevant, patient-oriented reasons may explain why patients discontinue treatments," the authors write. "These results may inform the development of [public policy](#) and evidence-based strategies to improve treatment satisfaction and to maintain successful long-term [psoriasis](#) control."

Several authors disclosed financial ties to the [pharmaceutical industry](#).

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Reasons for discontinuation vary by psoriasis treatment (2012, August 10) retrieved 31 March 2023 from
<https://medicalxpress.com/news/2012-08-discontinuation-vary-psoriasis-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--