

Urban HIV infection mainly due to male-male sexual contact

November 30 2012



More than half of HIV infections in metropolitan statistical areas, smaller metropolitan areas, and nonmetropolitan areas in the United States and Puerto Rico can be attributed to male-to-male sexual contact, according to research published in the Nov. 30 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

(HealthDay)—More than half of HIV infections in metropolitan statistical areas (MSAs), smaller metropolitan areas, and nonmetropolitan areas in the United States and Puerto Rico can be attributed to male-to-male sexual contact, according to research published in the Nov. 30 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

Hollie Clark, M.P.H., from the CDC in Atlanta, and colleagues estimated the number of HIV infections in newly diagnosed individuals in 2010 and classified them by transmission category and location. The authors focused on geographic differences in the prevalence of [HIV](#)

[infection](#) from male-to-male sexual contact among individuals age 13 and older in the United States and Puerto Rico.

The authors note that the largest percentage of HIV infections in MSAs, smaller metropolitan areas, and nonmetropolitan areas can be attributed to male-to-male sexual contact (62.1, 56.1, and 53.7 percent, respectively). Of the cases of HIV infection due to male-to-male sexual contact (28,851 cases), 81.7 percent were in MSAs, 48.4 percent of which resided in seven MSAs that represented 31.7 percent of the total population of individuals aged 13 years and older.

"The results of this analysis underscore the uneven geographic distribution of the burden of HIV infection in MSAs in the United States and Puerto Rico," write the authors of an editorial note. "The geographic disparity in HIV burden also indicates a need to target men who have sex with men who bear a large percentage of the burden of infection in areas where persons are at greatest risk for [HIV transmission](#)."

More information: [Full Text](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Urban HIV infection mainly due to male-male sexual contact (2012, November 30) retrieved 22 January 2023 from <https://medicalxpress.com/news/2012-11-urban-hiv-infection-due-male-male.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--