

180,000 deaths worldwide may be associated with sugary soft drinks

19 March 2013

Sugar-sweetened sodas, sports drinks and fruit drinks may be associated with about 180,000 deaths around the world each year, according to research presented at the American Heart Association's Epidemiology and Prevention/Nutrition, Physical Activity and Metabolism 2013 Scientific Sessions.

Sugar-sweetened beverages are consumed throughout the world, and contribute to [excess body weight](#), which increases the risk of developing diabetes, cardiovascular diseases and some cancers. Using data collected as part of the 2010 Global Burden of Diseases Study, the researchers linked intake of sugar-sweetened beverages to 133,000 diabetes deaths, 44,000 deaths from cardiovascular diseases and 6,000 cancer deaths. Seventy-eight percent of these deaths due to over-consuming [sugary drinks](#) were in low and middle-income countries, rather than high-income countries.

"In the U.S., our research shows that about 25,000 deaths in 2010 were linked to drinking sugar-sweetened beverages," said Gitanjali M. Singh, Ph.D., co-author of the study and a postdoctoral research fellow at the Harvard School of Public Health in Boston, Mass.

Researchers calculated the quantities of sugar-sweetened [beverage intake](#) around the world by age and sex; the effects of this consumption on obesity and diabetes; and the impact of obesity and diabetes-related deaths. Of nine world regions, Latin America/Caribbean had the most diabetes deaths (38,000) related to the consumption of sugar-sweetened beverages in 2010. East/Central Eurasia had the largest numbers of cardiovascular deaths (11,000) related to sugary [beverage consumption](#) in 2010. Among the world's 15 most populous countries, Mexico—one of the countries with the highest per-capita consumption of sugary beverages in the world—had the highest death rate due to these

beverages, with 318 deaths per million adults linked to sugar-sweetened beverage intake.

Japan, one of the countries with lowest per-capita consumption of sugary beverages in the world, had the lowest death rate associated with the consumption of sugary beverages, at about 10 deaths due to per million adults.

"Because we were focused on deaths due to chronic diseases, our study focused on adults. Future research should assess the amount of sugary beverage consumption in children across the world and how this affects their current and future health," Singh said.

The Global Burden of Disease Study 2010 is an international, collaborative, systematic effort to quantify the global distribution and causes of major diseases, injuries and health risk factors.

The [American Heart Association](#) recommends adults consume no more than 450 calories per week, from sugar-sweetened beverages, based on a 2,000 calorie diet and offers tips on how Life's Simple 7™ can help you make better lifestyle choices and eat healthier.

Provided by American Heart Association

APA citation: 180,000 deaths worldwide may be associated with sugary soft drinks (2013, March 19) retrieved 29 June 2022 from <https://medicalxpress.com/news/2013-03-deaths-worldwide-sugary-soft.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.