


Elderly benefit from using implantable defibrillators

June 17 2013



Young or old, implantable defibrillators can help.

- An implantable cardioverter defibrillator (ICD) is a small battery powered device placed under the skin of the chest which delivers electrical shocks to restore a normal heartbeat if it detects a dangerous abnormal rhythm.
- Older people are just as likely to get an appropriate shock as younger people.
- Overall health, not age alone, should help decide whether someone gets an ICD.

Source - American Heart Association journal *Circulation*; D. Lee

This infographic from the article outlines the elderly may benefit from implantable cardioverter defibrillators as much as younger people. Credit: American Heart Association journal *Circulation*; D. Lee

The elderly may benefit from implantable cardioverter defibrillators as much as younger people, according to new research in the American Heart Association journal *Circulation*.

An implantable cardioverter defibrillator (ICD) is a small battery-powered device placed under the skin of the chest which delivers [electrical impulses](#) to restore a normal heartbeat if it detects a dangerous abnormal rhythm.

Overall health—not age alone—should determine how well patients will do after getting an ICD and help guide decisions about who should receive one, researchers said.

"Whether elderly patients benefit from the devices has been controversial and research on the topic is lacking," said Douglas S. Lee, M.D., Ph.D., lead author and scientist at the Institute for Clinical Evaluative Sciences and cardiologist at the Peter Munk Cardiac Centre in Toronto, Ontario, Canada. "The issue is important as the population ages and the number of elderly people living with heart disease grows."

Researchers studied 5,399 patients who had ICDs implanted because of poor [heart function](#) due to a prior heart attack or [heart failure](#), or after being resuscitated from [cardiac arrest](#).

"Older patients were just as likely to experience an appropriate [electrical shock](#) from the device to treat a life-threatening heart rhythm. However, older patients experienced more non-cardiac and cardiovascular hospitalizations and higher associated rates of death overall," said Lee, who is also associate professor of medicine at the University of Toronto.

Other study findings include:

- Among those who received an ICD with heart failure or after

heart attack, 38.5 percent were 70 years or older and 7 percent were 80 years or older.

- Among those who received an ICD after surviving a cardiac arrest, 42.5 percent were 70 years or older and 10.9 percent were in their 80s.
- Older age correlated with increased risk of death after ICD implantation in both groups. For example, among those undergoing ICD implantation for prevention of cardiac arrest, 2 in every 100 patients ages 18-49 died whereas 10 in 100 died among those 80 years and older.
- Use of the devices was equally effective among younger and older patients in restoring heart rhythm after a serious life-threatening rhythm disorder occurred. Across all age groups, ICDs more often delivered appropriate electrical shocks when implanted after cardiac arrest than when used to prevent the first cardiac arrest.

Provided by American Heart Association

Citation: Elderly benefit from using implantable defibrillators (2013, June 17) retrieved 4 April 2023 from <https://medicalxpress.com/news/2013-06-elderly-benefit-implantable-defibrillators.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.