

In mild strokes, ultra-early treatment may eliminate risk of disability

22 August 2013

In the case of mild or moderate strokes, getting treatment ultra-fast – within 90 minutes of experiencing symptoms – greatly reduces the risk of suffering disability, according to a new study reported in the American Heart Association's journal *Stroke*.

The American Heart Association/American Stroke Association recommends getting to a hospital within three hours of the onset of [stroke symptoms](#). According to guidelines, clot-busting drugs may be given to treat stroke up to 4.5 hours after the onset of symptoms.

The study found that survivors with mild to moderate strokes who were given the clot-busting drugs in the first 90 minutes of the recommended [time window](#) had little or no disability three months later compared to those who were treated between 90 and 270 minutes.

"Ultra-early treatment increases the likelihood of excellent outcome in patients with moderately severe symptoms, and in secondary analysis also in those with mild symptoms," said Daniel Strbian, M.D., Ph.D., from the Department of Neurology at Helsinki University Central Hospital in Helsinki, Finland. "All measures must be taken to reduce onset-to-treatment time as much as possible."

The study included more than 6,800 [stroke patients](#) at 10 [stroke centers](#) in Europe over 14 years. They were treated intravenously with Alteplase, a clot-busting drug that is given through an IV in the vein. Patients were separated into three groups based on stroke severity – minor (NIH stroke score 0-6), mild/moderate (NIH score 7-12), or moderate/severe (NIH score higher than 12). Those with mild to moderate stroke seemed to benefit most from the ultra-early care. Early treatment also helped those with minor strokes, but the likelihood of disability is already very low in these patients.

Those with severe stroke did not benefit as much from the ultra-early treatment because they had severe [artery blockage](#).

"We need more research to offer something more for people with severe strokes," Strbian said.

Provided by American Heart Association

APA citation: In mild strokes, ultra-early treatment may eliminate risk of disability (2013, August 22) retrieved 25 June 2022 from <https://medicalxpress.com/news/2013-08-mild-ultra-early-treatment-disability.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.