

Bipolar drugs lose effect during pregnancy, so women need higher doses to stay well

November 1 2013

New Northwestern Medicine research offers one of the first in-depth studies of how physiological changes during pregnancy reduce the effects of a commonly used drug to treat bipolar disorder, making women more vulnerable to recurring episodes. The new findings will help psychiatrists and physicians prevent bipolar manic and depressive symptoms during pregnancy, which are risky for the health of the mother and her unborn child.

When a woman with bipolar disorder becomes pregnant, she and her physician often don't realize her medication needs adjusting to prevent the symptoms from coming back – a higher risk during pregnancy. There also is little information and research to guide dosing for psychiatric medications during pregnancy.

Approximately 4.4 million women in the U.S. have bipolar disorder with women of childbearing age having the highest prevalence.

The new study shows the blood concentration of the commonly used drug lamotrigine decreases in [pregnant women](#). About half of the women in the study had worsening [depressive symptoms](#) as their lamotrigine blood levels dropped. The drug levels fall because women have increased metabolism during pregnancy.

"Now physicians change the dose of the drug in response to women's symptoms worsening," said lead investigator Crystal Clark, M.D., an assistant professor of psychiatry and behavioral sciences at Northwestern

University Feinberg School of Medicine and a psychiatrist at Northwestern Memorial Hospital. "We need to optimize their medication dosing so they stay well."

The study results will help physicians understand how to increase their patients' doses during [pregnancy](#) and then reduce them postpartum to avoid toxicity, Clark said. Guidelines for prescribing the [drug](#) for pregnant women with the disorder also are included.

The study was published Nov. 1 in the *American Journal of Psychiatry*.

Depressive episodes—as opposed to manic—are most likely to recur in pregnant women with [bipolar disorder](#).

"The safety of the fetus is at risk," Clark said. "Pregnant women that are depressed are less likely to take care of themselves which often leads to poor nutrition, lack of compliance with prenatal care and isolation from family and friends. It has also been linked to premature births and babies with low birth weights among other poor birth outcomes."

Provided by Northwestern University

Citation: Bipolar drugs lose effect during pregnancy, so women need higher doses to stay well (2013, November 1) retrieved 2 April 2023 from <https://medicalxpress.com/news/2013-11-bipolar-drugs-effect-pregnancy-women.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--