


Cell-saving drugs could reduce brain damage after stroke

March 26 2014



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

Long-term brain damage caused by stroke could be reduced by saving cells called pericytes that control blood flow in capillaries, reports a new study led by scientists from University College London.

Until now, many scientists believed that blood flow within the brain was solely controlled by changes in the diameter of arterioles, [blood vessels](#)

that branch out from arteries into smaller capillaries. The latest research reveals that the brain's blood supply is in fact chiefly controlled by the narrowing or widening of capillaries as pericytes tighten or loosen around them.

The study, published this week in *Nature*, shows not only that pericytes are the main regulator of blood flow to the brain, but also that they tighten and die around [capillaries](#) after stroke. This significantly impairs blood flow in the long term, causing lasting damage to brain cells. The team of scientists from UCL, Oxford University and the University of Copenhagen showed that certain chemicals could halve pericyte death from simulated stroke in the lab, and hope to develop these into drugs to treat [stroke victims](#).

"At present, clinicians can remove clots blocking blood flow to the brain if [stroke patients](#) reach hospital early enough," explains Professor David Attwell of UCL's Department of Neuroscience, Physiology & Pharmacology, who led the study. "However, the capillary constriction produced by pericytes may, by restricting the blood supply for a long time, cause further damage to nerve cells even after the clot is removed. Our latest research suggests that devising drugs to prevent capillary constriction may offer new therapies for reducing the disability caused by stroke."

"This discovery offers radically new treatment approaches for stroke," says study co-author Professor Alastair Buchan, Dean of Medicine and Head of the Medical Sciences Division at Oxford University.

"Importantly, we should now be able to identify drugs that target these cells. If we are able to prevent pericytes from dying, it should help restore blood flow in the brain to normal and prevent the ongoing slow damage we see after a [stroke](#) which causes so much neurological disability in our patients."

The new research also gives insight into the mechanisms underlying the

use of functional magnetic resonance imaging to detect blood flow changes in the brain.

"Functional imaging allows us to see the activity of nerve cells within the human [brain](#) but until now we didn't quite know what we were looking at," explains Professor Attwell. "We have shown that pericytes initiate the increase in [blood flow](#) seen when [nerve cells](#) become active, so we now know that functional imaging signals are caused by a pericyte-mediated increase of capillary diameter. Knowing exactly what functional imaging shows will help us to better understand and interpret what we see."

More information: Capillary pericytes regulate cerebral blood flow in health and disease, [DOI: 10.1038/nature13165](https://doi.org/10.1038/nature13165)

Provided by University College London

APA citation: Cell-saving drugs could reduce brain damage after stroke (2014, March 26) retrieved 13 December 2022 from <https://medicalxpress.com/news/2014-03-cell-saving-drugs-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--