

Patient variables predict expectations in spine surgery

22 June 2014



"Multiple clinical variables were associated with expectations, with younger age and more disability due to pain being the most consistently associated with more expectations," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

(HealthDay)—Clinical and demographic variables can predict patients' expectations regarding anticipated improvement from cervical spine surgery, according to a study published June 15 in *Spine*.

Carol A. Mancuso, M.D., from the Hospital for Special Surgery in New York, and colleagues analyzed surveys completed by 150 [patients](#) (mean age 54 years; 61 percent male) several days before cervical surgery. The Expectations Survey comprises 20 physical and psychological items. Disability due to pain was evaluated with the Neck Disability Index (NDI), and overall physical and mental health was measured with the 12-item Short Form Health Survey.

The researchers found that the most commonly chosen items were relieve neck (87 percent) and upper extremity (85 percent) pain, stop the [spine](#) condition from getting worse (97 percent), and remove the control the spine condition had on life (96 percent). Patients were more likely to choose more items if they were younger (odds ratio [OR], 2.2; 95 percent confidence interval [CI], 1.2 to 4.0; P = 0.01), had worse NDI scores (OR, 6.5; 95 percent CI, 3.2 to 13.2; P

APA citation: Patient variables predict expectations in spine surgery (2014, June 22) retrieved 27 October 2022 from <https://medicalxpress.com/news/2014-06-patient-variables-spine-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.