


# Physician, system factors affect CAD detection rates

September 11 2014

---



(HealthDay)—Nonclinical factors account for considerable variation in the detection of obstructive coronary artery disease (CAD) with coronary angiogram, according to a study published online Sept. 2 in *Circulation: Cardiovascular Quality & Outcomes*.

Harindra C. Wijeyesundera, M.D., Ph.D., from the University of Toronto in Canada, and colleagues evaluated data from all [patients](#) in Ontario, Canada who underwent [coronary angiography](#) for possible suspected stable ischemic heart disease from Oct. 1, 2008 to Sept. 30, 2011. The incremental impact of nonclinical factors on variation in detection of obstructive CAD was assessed, with physician self-referral for angiography, physician type (invasive or interventional), and hospital type examined as the nonclinical factors of interest.

The researchers found that 52 percent of the 60,986 patients had obstructive CAD, with significant range across hospitals (37.3 to 69.2 percent). Fewer cases of obstructive CAD were seen in self-referral versus nonself-referral patients (49.8 versus 53.5 percent; odds ratio [OR], 0.89; P<0.05). "Physician and system factors are important predictors of obstructive CAD with coronary angiography," conclude the authors.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Physician, system factors affect CAD detection rates (2014, September 11) retrieved 21 February 2023 from  
<https://medicalxpress.com/news/2014-09-physician-factors-affect-cad.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--