

# Depression increases risk of falls in elderly

October 7 2014, by Siobhan Moylan

---



Depressive symptoms, antidepressant use, poorer balance and poorer cognitive functioning are all associated with increased risk of falls in the elderly. Credit: Thinkstock

Falls are common in elderly people but the risk increases markedly when they have depressive symptoms, research from Neuroscience Research Australia shows.

Depressive symptoms, antidepressant use, poorer balance and poorer cognitive functioning are all associated with increased [risk](#) of [falls](#) in the elderly. The researchers investigated the interactions between and the significance of these [risk factors](#).

"Risk of falls increased by 55 per cent in participants with any two risk factors and by 144 per cent in participants with three to four risk factors," says study co-author Dr Kim Delbaere, from NeuRA.

"The more risk factors a participant had, the more likely they were to experience a fall," says Dr Delbaere, also a conjoint senior lecturer at UNSW.

Depression and use of antidepressants increased the risk of falls independently of reduced cognitive and physical functioning.

"Depressive symptoms are a potentially modifiable risk factor for falls and should therefore be addressed in fall prevention programs, especially in high risk populations,"

The researchers suggest that therapies to reduce the likelihood of falls among older people experiencing depression should focus on a combination of physical exercise, including resistance and balance training, and psychosocial therapies such as mindfulness.

"Given that antidepressant medication may increase fall risk, treatment of [depressive symptoms](#) using non-pharmacological approaches should be considered as part of all [prevention programs](#), especially in populations at high risk of falls," Dr Delbaere says. "These could include stress reduction and relaxation techniques, and other cognitive behavioural and mindfulness techniques."

The study relied on data from 488 people aged 70 to 90 living in the community in eastern Sydney who are taking part in the Sydney Memory & Ageing Study being conducted by UNSW's CHeBA. The participants underwent comprehensive psychological, cognitive and physiological assessment, and were monitored for falls over the course of a year.

**More information:** "Depressive symptoms increase fall risk in older people, independent of antidepressant use, and reduced executive and physical functioning." Tasha Kvelde, et al. *Archives of Gerontology and Geriatrics* April 28, 2014 DOI: [dx.doi.org/10.1016/j.archger.2014.09.003](https://doi.org/10.1016/j.archger.2014.09.003)

Provided by University of New South Wales

Citation: Depression increases risk of falls in elderly (2014, October 7) retrieved 2 December 2023 from <https://medicalxpress.com/news/2014-10-depression-falls-elderly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.