

Ankle, knee strength generation slower with diabetic neuropathy

October 29 2014



(HealthDay)—When walking up and down stairs, patients with diabetic peripheral neuropathy (DPN) are slower at generating strength at the ankle and knee compared to control participants, which may increase the risk of falls, according to a study published online Oct. 14 in *Diabetes Care*.

Joseph C. Handsaker, from the Manchester Metropolitan University in the United Kingdom, and colleagues examined 63 participants (21 patients with DPN, 21 controls with diabetes, and 21 healthy controls) walking up and down a custom-built staircase. Analysis included assessment of speed of strength generation at the ankle and [knee](#) and muscle activation patterns of the ankle and knee extensor muscles.

The researchers found that patients with neuropathy displayed

significantly slower ankle and knee strength generation than healthy controls during stair ascent and descent (P ankle and knee extensors took significantly longer to reach peak activation (P

"These changes, which are likely caused by altered activations of the extensor muscles, increase the likelihood of instability and may be important contributory factors for the increased risk of falling," the authors write. "Resistance exercise training may be a potential clinical intervention for improving these aspects and thereby potentially reducing fall risk."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Ankle, knee strength generation slower with diabetic neuropathy (2014, October 29)
retrieved 23 April 2023 from
<https://medicalxpress.com/news/2014-10-ankle-knee-strength-slower-diabetic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--