

Spinal surgery varies by region in the US

November 3 2014

(HealthDay)—Surgery for low back pain caused by spinal stenosis varies depending on where in the United States you live, according to a Dartmouth Atlas Project report.

The analysis of Medicare data revealed that rates of spinal decompression varied eight-fold across the United States, from 25.3 procedures per 100,000 patients in Bronx, N.Y., to 216.7 procedures per 100,000 patients in Mason City, Iowa. In general, rates of spinal decompression were highest in the Pacific Northwest and northern Mountain states.

Rates of spinal fusion varied more than 14-fold nationally, from 9.2 procedures per 100,000 patients in Bangor, Maine, to 127.5 procedures per 100,000 patients in Bradenton, Fla., according to the report.

"Nearly 80 percent of Americans will experience [low back pain](#) at some point in their lives, and about 30 million people a year receive professional medical care for a spine problem," coauthor Brook Martin, Ph.D., M.P.H., of the Dartmouth Institute of Health Policy & Clinical Practice in Lebanon, N.H., said in a college news release.

More information: [Full Article](#)
[More Information](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

APA citation: Spinal surgery varies by region in the US (2014, November 3) retrieved 8 December 2022 from <https://medicalxpress.com/news/2014-11-spinal-surgery-varies-region.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.