

# Omega-3 PUFAs can reduce smoking, cut tobacco craving

13 November 2014



Further studies are needed on larger samples to explore the possible therapeutic implications for heavy cigarette smokers."

Solgar Vitamin and Herb Company donated the omega-3 capsules used in the study.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

(HealthDay)—Supplementation with omega-3 polyunsaturated fatty acids (PUFAs) can reduce daily smoking and tobacco craving, according to a study published in the *Journal of Psychopharmacology*.

Noting that low concentrations of omega-3 PUFAs can affect neurotransmission and may increase cigarette cravings, Sharon Rabinovitz, M.D., from the University of Haifa in Israel, examined the effect of PUFA supplementation on tobacco craving. Forty-eight regular [cigarette smokers](#) were administered 2,710 mg eicosapentaenoic acid/day and 2,040 mg docosahexaenoic acid/day for one month in a pilot study.

Rabinovitz found that PUFA supplementation for one month correlated with a significant reduction in reported daily smoking and in tobacco craving following cigarette cue exposure. In the month following treatment discontinuation, craving did not return to baseline values.

"This is the first study demonstrating that omega-3 PUFA supplementation reduces tobacco craving in regular smokers, compared to placebo treatment," Rabinovitz writes. "Thus, omega-3 PUFAs may be of benefit in managing [tobacco consumption](#)."

APA citation: Omega-3 PUFAs can reduce smoking, cut tobacco craving (2014, November 13) retrieved 12 June 2022 from <https://medicalxpress.com/news/2014-11-omega-pufas-tobacco-craving.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*