


Early retina cell changes in glaucoma identified

February 11 2015



An example of retinal ganglion cells with dendrites in the retina of a healthy eye.
Credit: UC San Diego School of Medicine

Glaucoma, the second leading cause of blindness, usually stems from elevated eye pressure, which in turn damages and destroys specialized neurons in the eye known as retinal ganglion cells. To better understand these cellular changes and how they influence the progression and severity of glaucoma, researchers at University of California, San Diego

School of Medicine and Shiley Eye Institute turned to a mouse model of the disease. Their study, published Feb. 10 in *The Journal of Neuroscience*, reveals how some types of retinal ganglion cells alter their structures within seven days of elevated eye pressure, while others do not.

"Understanding the timing and pattern of cellular changes leading to retinal ganglion cell death in glaucoma should facilitate the development of tools to detect and slow or stop those [cellular changes](#), and ultimately preserve vision," said Andrew D. Huberman, PhD, assistant professor of neurosciences, neurobiology and ophthalmology. Huberman co-authored the study with Rana N. El-Danaf, PhD, a postdoctoral researcher in his lab.

Retinal ganglion cells are specialized neurons that send visual information from the eye's retina to the brain. Increased pressure within the eye can contribute to retinal ganglion cell damage, leading to glaucoma. Even with pressure-lowering drugs, these cells eventually die, leading to vision loss.

In this study, Huberman and El-Danaf used a [mouse model](#) engineered to express a green fluorescent protein in specific retinal ganglion cell subtypes. This tool allowed them to examine four subtypes of [retinal ganglion](#) cells. The different cell types differ by the location in the eye to which they send the majority of their dendrites (cellular branches). Within seven days of elevated eye pressure, all [retinal ganglion cells](#) that send most or all of their dendrites to a region of the eye known as the OFF sublamina underwent significant rearrangements, such as reductions in number and length of dendritic branches. Retinal [ganglion cells](#) with connections in the ON part of the retina did not.

"We are very excited about this discovery," Huberman said. "One of the major challenges to the detection and treatment of glaucoma is that you

have to lose a lot of cells or [eye pressure](#) has to go way up before you know you have the disease. These results tell us we should design visual field tests that specifically probe the function of certain retinal cells. In collaboration with the other researcher members of the Glaucoma Research Foundation Catalyst for a Cure, we are doing just that and we are confident these results will positively impact human patients in the near-future."

More information: *The Journal of Neuroscience*,
www.jneurosci.org/content/35/6/2329.short

Provided by University of California - San Diego

Citation: Early retina cell changes in glaucoma identified (2015, February 11) retrieved 15 February 2024 from <https://medicalxpress.com/news/2015-02-early-retina-cell-glaucoma.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--