

Study shows long-term effects of type 2 diabetes on the brain, thinking

8 July 2015

In just two years, people with type 2 diabetes experienced negative changes in their ability to regulate blood flow in the brain, which was associated with lower scores on tests of cognition skills and their ability to perform their daily activities, according to a new study published in the July 8, 2015, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

"Normal [blood flow](#) regulation allows the brain to redistribute blood to areas of the brain that have increased activity while performing certain tasks," said study author Vera Novak, MD, PhD, of Harvard Medical School in Boston. "People with [type 2 diabetes](#) have impaired blood flow regulation. Our results suggest that diabetes and [high blood sugar](#) impose a chronic negative effect on cognitive and decision-making skills."

The study involved 40 people with an average age of 66. Of those, 19 had type 2 diabetes and 21 did not have diabetes. Those with diabetes had been treated for the disease for an average of 13 years. The participants were tested at the beginning of the study and again two years later. Tests included cognition and memory tests, MRI scans of the brain to look at brain volume and blood flow, and blood tests to measure control of blood sugar and inflammation.

After two years, the people with diabetes had decreases in their ability to regulate blood flow in the brain. They also had lower scores on several tests of memory and thinking skills. People with lower ability to regulate blood flow at the beginning of the study had greater declines in a measure of how well they could complete [daily activities](#) such as bathing and cooking.

Higher levels of inflammation were also associated with greater decreases in blood flow regulation, even if people had good control of their diabetes and blood pressure, Novak said.

On a test of learning and memory, the scores of the people with diabetes decreased by 12 percent, from 46 points to 41 points over the two years of the study, while the scores of those without diabetes stayed the same, at 55 points. Blood flow regulation in the [brain](#) was decreased by 65 percent in people with [diabetes](#).

"Early detection and monitoring of blood flow regulation may be an important predictor of accelerated changes in cognitive and decision-making skills," Novak said.

She said additional studies involving more people and extending for a longer time period are needed to better understand the relationship and timing with [blood flow regulation](#) and changes in thinking and memory skills.

Provided by American Academy of Neurology

APA citation: Study shows long-term effects of type 2 diabetes on the brain, thinking (2015, July 8) retrieved 10 September 2022 from <https://medicalxpress.com/news/2015-07-long-term-effects-diabetes-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.