

Metformin linked to lower risk of head, neck cancer

9 September 2015





(HealthDay)—Metformin is associated with a lower risk of developing head and neck cancer in patients with diabetes, according to a study published in the September issue of *Head & Neck*.

Yung-Chang Yen, M.D., Ph.D., from the Chi Mei Medical Centre in Tainan City, Taiwan, and colleagues compared the risk of head and neck cancer in 2011 among 66,600 patients, all newly diagnosed with diabetes in 2002. Metformin was subsequently taken by half of the patients. The two groups were matched for comorbidities, sex, and age.

The researchers found that the incidence of head and neck cancer was 34 percent lower in the metformin group than in the control group (adjusted hazard ratio [HR], 0.66). Similarly, among those taking metformin, the risks for oropharyngeal cancer (adjusted HR, 0.66) and nasopharyngeal carcinoma (adjusted HR, 0.5) were significantly lower.

"These findings indicate the potential clinical benefits of using metformin to prevent patients with diabetes from developing head and neck cancer," the authors write.

More information: Abstract

Full Text (subscription or payment may be required)



APA citation: Metformin linked to lower risk of head, neck cancer (2015, September 9) retrieved 29 September 2022 from https://medicalxpress.com/news/2015-09-metformin-linked-neck-cancer.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.