

Modest predictive power for HbA1c in atherosclerotic CVD risk

September 10 2015



(HealthDay)—In the context of conventional cardiovascular disease (CVD) risk factors, hemoglobin A1c (HbA1c) has a modest effect on predicted atherosclerotic CVD risk, according to a study published online Sept. 8 in *Circulation: Cardiovascular Quality and Outcomes*.

Jamie A. Jarmul, from the University of North Carolina at Chapel Hill, and colleagues examined the effect of HbA1c on CVD risk in the context of other CVD <u>risk factors</u>. HbA1c and other CVD risk factor measurements were analyzed in 2,000 individuals aged 40 to 79 years without preexisting diabetes mellitus or CVD. For a set of example scenarios, post-test 10-year atherosclerotic CVD risk was calculated incorporating the actual versus predicted HbA1c, according to established methods.



The researchers found that significant predictors in the model included age, sex, race/ethnicity, and traditional <u>cardiovascular risk factors</u>, with the expected HbA1c distribution significantly higher in non-Hispanic black, non-Hispanic Asian, and Hispanic individuals versus non-Hispanic whites/others. A modest effect was seen on post-test atherosclerotic CVD risk for incorporation of the expected HbA1c distribution into pretest atherosclerotic CVD risk. In the patient examples, depending on the scenario, HbA1c of

Citation: Modest predictive power for HbA1c in atherosclerotic CVD risk (2015, September 10) retrieved 12 March 2023 from <u>https://medicalxpress.com/news/2015-09-modest-power-hba1c-atherosclerotic-cvd.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.