

Risk of fractures reduced in polycystic ovary syndrome

16 November 2015



"Women with PCOS had reduced risk of fractures, in particular of the appendicular skeleton," the authors write. "The risk reduction was greater in women with younger age at diagnosis, suggesting that the skeletal effects of PCOS may be greater in women who have not yet reached peak bone mass."

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2015 HealthDay. All rights reserved.

(HealthDay)—The risk of fractures is reduced in women with polycystic ovary syndrome (PCOS), according to a study published online Nov. 6 in the *Journal of Bone and Mineral Research*.

Katrine Hass Rubin, Ph.D., from the University of Southern Denmark in Odense, and colleagues examined <u>fracture risk</u> in patients with PCOS. Women with PCOS and/or hirsutism were identified from the Danish National Patient Register (19,199 women), and assigned three age-matched controls (57,483 women) on the index date of diagnosis of PCOS.

The researchers found that within the cohort. fracture rates were decreased for women with PCOS versus controls (10.3 versus 13.6/1,000 patient-years) The adjusted odds ratios were 0.76, 0.82, and 0.57 for all <u>fractures</u>, major osteoporotic fractures, and fractures of the head and face, respectively. Women diagnosed before age 30 years had more pronounced risk reduction. Women with PCOS had significantly more strainand sprain-related hospital contact. Within a wellcharacterized subcohort of 1,217 women, the researchers observed no difference in the risk reduction in fractures for PCOS women with normal versus elevated testosterone levels; risk reduction was slightly smaller in overweight women.



APA citation: Risk of fractures reduced in polycystic ovary syndrome (2015, November 16) retrieved 11 September 2022 from https://medicalxpress.com/news/2015-11-fractures-polycystic-ovary-syndrome.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.