

Increased long-term death risk for adolescents hospitalized for adversity-related injury

29 December 2015

Adolescents discharged from hospitals in England after an admission for violent, drug- or alcohol-related, or self-inflicted injuries have increased risks of subsequent death and emergency re-admission up to a decade later, according to a study published this week in *PLOS Medicine*. The study, conducted by Annie Herbert at University College London, UK, and colleagues, showed that in England, risks of death after all types of adversity-related injury were higher than after accident-related injury (61% (95% CI 43%-82%) higher in girls and 113% (95% CI 98%-129%) higher in boys).

Adolescents who present with an adversity-related injury often re-present later with other adversity-related injuries. However, national guidance in England stipulates psychosocial assessment only for presentations of self-inflicted injury. To determine which adolescents in the broader group are at elevated risk of further harm, Herbert and [colleagues](#) used National Health Service hospital admissions data from 1997 into 2012 for 10-19 year olds with emergency admissions for adversity-related or accident-related injury (333,009 and 649,818 adolescents, respectively). Among adolescents discharged after adversity-related injury, one in 137 girls and one in 64 boys died within ten years, and 54.2% of girls and 40.5% of boys had a subsequent emergency re-admission. These rates were roughly 1.5-2 times higher than after accident-related injury. Risks of death were highest in 18-19 year olds (one in 52 boys and one in 90 girls), and those with either chronic conditions (typically mental/behavioural or respiratory disorders for [adolescents](#) in this study) or who lived in deprived areas.

Misclassification of some adversity-related injuries as accident-related injuries, and residual confounding by unknown characteristics shared

within the two groups, may affect the accuracy of these findings. Nevertheless, these findings identify a broader range of risk factors for subsequent harm. The authors state, "[t]hese findings justify extending national policy for psychosocial assessment after self-inflicted injury to all types of adversity-related injury."

More information: Annie Herbert et al. 10-y Risks of Death and Emergency Re-admission in Adolescents Hospitalised with Violent, Drug- or Alcohol-Related, or Self-Inflicted Injury: A Population-Based Cohort Study, *PLOS Medicine* (2015). [DOI: 10.1371/journal.pmed.1001931](https://doi.org/10.1371/journal.pmed.1001931)

Provided by Public Library of Science

APA citation: Increased long-term death risk for adolescents hospitalized for adversity-related injury (2015, December 29) retrieved 16 September 2022 from <https://medicalxpress.com/news/2015-12-long-term-death-adolescents-hospitalized-adversity-related.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.