

Current malaria treatment fails in Cambodia due to drug-resistant parasites

7 January 2016



Credit: CDC

New findings from the National Institute of Allergy and Infectious Diseases (NIAID), part of the National Institutes of Health (NIH), confirm dihydroartemisinin-piperaquine, the first-line treatment for *Plasmodium falciparum* malaria infection in Cambodia, has failed in certain provinces due to parasite resistance to artemisinin and piperaquine. Dihydroartemisinin-piperaquine is an artemisinin combination therapy (ACT) for malaria that combines potent, fast-acting artemisinin with a long-acting partner drug, piperaquine. Resistance to artemisinin in parts of Southeast Asia is [well-documented](#), but until now only a few studies have presented clear evidence of piperaquine resistance. Additional study findings suggest that artesunate, a form of artemisinin, plus mefloquine, a different long-acting partner drug, should be the first-line ACT in areas where dihydroartemisinin-piperaquine treatment has failed, the study authors note.

NIAID researchers and colleagues sought to confirm the presence of piperaquine-resistant infections in Cambodia by comparing the efficacy of dihydroartemisinin-piperaquine treatment in 204

malaria-afflicted [participants](#) aged 2 to 65 years from three provinces in Cambodia with varying levels of artemisinin [resistance](#). After monitoring parasite levels in the blood for 63 days, investigators found parasites had reemerged despite initial clearance in 45.7 percent of participants in Pursat, 15.9 percent of participants in Preah Vihear and 1.67 percent of participants in Ratanakiri. The results indicate the ACT is failing in Pursat and Preah Vihear, where artemisinin resistance is common, but remains highly efficacious in Ratanakiri, where resistance is uncommon.

Laboratory tests showed the parasites from dihydroartemisinin-piperaquine failures contained a genetic marker of artemisinin resistance and had a decreased susceptibility to piperaquine, demonstrating that both artemisinin and piperaquine resistance contributed to treatment failures. However, the parasites also showed an increased susceptibility to mefloquine and completely lacked the molecular marker for mefloquine resistance. These findings informed new WHO guidelines reinstating artesunate plus mefloquine as the first-line ACT in Cambodia where dihydroartemisinin-piperaquine [treatment](#) has failed. The findings also provide evidence to initiate surveillance programs to track the spread of piperaquine resistance and clinical trials to test alternative combination therapies.

More information: Amaratunga C, Lim P, et al. Dihydroartemisinin-piperaquine resistance in *Plasmodium falciparum* malaria in Cambodia: a multi-site observational cohort study. *The Lancet Infectious Diseases* [DOI: 10.1016/S1473-3099\(15\)00487-9](#) (2016).

Provided by NIH/National Institute of Allergy and Infectious Diseases

APA citation: Current malaria treatment fails in Cambodia due to drug-resistant parasites (2016, January 7) retrieved 3 December 2022 from <https://medicalxpress.com/news/2016-01-current-malaria-treatment-cambodia-due.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.