

Pre-pregnancy potato consumption may be linked to gestational diabetes risk

January 12 2016



Source: Wikipedia

Women who eat more potatoes before pregnancy may have higher rates of gestational diabetes—the form that occurs during pregnancy—compared to women who consume fewer potatoes, suggests a National Institutes of Health (NIH) study. The researchers propose that substituting potatoes with other vegetables, legumes or whole grains may help lower gestational diabetes risk. The findings appear in *The BMJ*.

Gestational diabetes is a common pregnancy complication that causes high [blood sugar levels](#) in the mother. The disorder can lead to future health problems for mother and child. Previous studies have linked foods with a [high glycemic index](#), a measure of the ability to raise blood sugar levels, to a higher risk of gestational or type 2 diabetes. However, until

the current study, the effect of potatoes, a common, high-glycemic food, on the development of gestational diabetes was unknown.

Researchers from NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) and Harvard University evaluated more than 15,000 women in the Nurses' Health Study II. They analyzed records from 1991 to 2001 of women who had no history of illness before pregnancy and had not had gestational diabetes before. Every four years, the women filled out a questionnaire on the kinds of foods they had eaten during the previous year. For potatoes, the women were asked if they had consumed baked, boiled, or mashed potatoes, fries or potato chips, with possible responses ranging from "never" to "six or more times a day."

The researchers found that [women](#) who ate more potatoes had a higher risk of gestational diabetes. They estimated reductions in gestational diabetes risk by substituting the following foods for two servings of [potatoes](#) per week:

- 9 percent for other vegetables
- 10 percent for legumes
- 12 percent for whole grain foods

The authors cautioned, however, that because their study was not designed to prove cause and effect, their results do not prove conclusively that potato consumption directly leads to [gestational diabetes](#). They added that their results need to be confirmed by other studies.

More information: Bao W, Tobias DK, Hu FB, Chavarro JE, and Zhang C. Pre-pregnancy potato consumption and risk of gestational diabetes mellitus: prospective cohort study. *BMJ* (2016) [dx.doi.org/10.1136/bmj.h6898](https://doi.org/10.1136/bmj.h6898)

Provided by National Institutes of Health

Citation: Pre-pregnancy potato consumption may be linked to gestational diabetes risk (2016, January 12) retrieved 22 January 2023 from <https://medicalxpress.com/news/2016-01-pre-pregnancy-potato-consumption-linked-gestational.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.