

# Liver recovery difficult even with improved diet—but faster if sugar intake is low

January 13 2016

---



Dietary sugar intake must come down, along with other improvements in diet and exercise, if there's a need to help repair liver damage, new research shows. Credit: Oregon State University

Liver damage caused by the typical "Western diet" - one high in fat, sugar and cholesterol that's common in developed countries such as the United States - may be difficult to reverse even if diet is generally improved, a new study shows.

The research, published today in *PLOS ONE* by scientists from Oregon State University, found that a diet with reduced fat and cholesterol helped, but did not fully resolve [liver damage](#) that had already been done - damage that in turn can lead to more serious health problems, such as cirrhosis or even cancer.

This study, done with laboratory animals, showed that diets low in fat and cholesterol could in fact aid with [weight loss](#), improved metabolism and health. But even then, if the diet was still high in sugar there was much less liver recovery, the scientists concluded.

The findings are significant, scientists say, because [liver problems](#) such as nonalcoholic fatty liver disease are surging in the U.S., affecting 10-35 percent of adults and an increasing number of children. The incidence of this problem can reach more than 60 percent in obese and type-2 diabetic populations.

"Many people eating a common American diet are developing extensive hepatic fibrosis, or scarring of their liver, which can reduce its capacity to function, and sometimes lead to cancer," said Donald Jump, a professor in the OSU College of Public Health and Human Sciences, principal investigator with the Linus Pauling Institute, and corresponding author on this research.

"There's a lot of interest in finding ways to help the liver recover from this damage, but this research suggests that diets lower in fat and cholesterol, even if they help you lose weight, are not enough," Jump said. "For more significant liver recovery, the intake of sugar has to

come down, probably along with other improvements in diet and exercise."

The issues are both serious and complex, the researchers said.

"Everyone recognizes this is a serious problem," said Kelli Lytle, an OSU doctoral candidate and lead author on this study. "We're trying to find out if some of the types of dietary manipulation that people use, such as weight loss based on a [low fat diet](#), will help address it. However, a common concern is that many 'low-fat' food products have higher levels of sugar to help make them taste better."

Weight loss does appear to help address some of the problems associated with the Western diet, the research shows. But according to this study, a diet with continued high levels of sugar will significantly slow recovery of liver damage that has already been done.

Complications related to liver inflammation, scarring and damage are projected to be the leading cause of [liver transplants](#) by 2020, the researchers noted in their study. Such scarring was once thought to be irreversible, but more recent research has shown it can be at least partially reversed with optimal diet and when the stimulus for liver injury is removed.

In this report, scientists studied two groups of laboratory mice that had been fed a "Western diet" and then switched to different, healthier diets, low in fat and cholesterol.

Both of the improved diets caused health improvements and weight loss. But one group that was fed a diet still fairly high in sugar - an amount of sugar comparable to the Western diet - had significantly higher levels of inflammation, oxidative stress and [liver fibrosis](#).

More research is still needed to determine whether a comprehensive program of [diet](#), weight maintenance, exercise and targeted drug therapies can fully resolve [liver](#) fibrosis, the study concluded.

**More information:** Kelli A. Lytle et al. Is Western Diet-Induced Nonalcoholic Steatohepatitis in Ldlr<sup>-/-</sup> Mice Reversible?, *PLOS ONE* (2016). [DOI: 10.1371/journal.pone.0146942](https://doi.org/10.1371/journal.pone.0146942)

Provided by Oregon State University

Citation: Liver recovery difficult even with improved diet—but faster if sugar intake is low (2016, January 13) retrieved 3 February 2024 from <https://medicalxpress.com/news/2016-01-liver-recovery-difficult-dietbut-faster.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.