

CDC: fewer blacks consistently retained in HIV care

February 8 2016



(HealthDay)—Fewer blacks are consistently retained in HIV care

compared with other racial/ethnic groups, according to research published in the Feb. 5 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

Sharoda Dasgupta, Ph.D., from the CDC in Atlanta, and colleagues used National HIV Surveillance System data to monitor progress toward reaching the National HIV/AIDS Strategy goals to improve care among persons living with HIV. Data were used to describe retention in care over three years and assess [differences](#) by race/ethnicity.

The researchers found that 38 percent of blacks with HIV infection were consistently retained in care during 2011 to 2013, compared with 50 and 49 percent of Hispanic/Latinos and non-Hispanic whites. When groups were stratified by sex or transmission category, the differences in consistent retention in care by race/ethnicity persisted. Among blacks, 35 and 44 percent of males and females, respectively, were consistently retained in care. Race/ethnicity-related differences in HIV care retention were established during the first year after diagnosis.

"Efforts to establish early HIV care among blacks are needed to mitigate racial/ethnic disparities in HIV outcomes over time," the authors write.

More information: [Full Text](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: CDC: fewer blacks consistently retained in HIV care (2016, February 8) retrieved 22 February 2023 from <https://medicalxpress.com/news/2016-02-cdc-blacks-retained-hiv.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--