

# Low-normal sodium deemed major risk for mortality in elderly

March 25 2016

---



A slightly lower serum sodium concentration within the normal range is

a major risk factor for mortality in elderly adults, according to a study published in the March issue of the *Journal of the American Geriatrics Society*.

Shin Y. Ahn, M.D., from Seoul National University Bundang Hospital in South Korea, and colleagues randomly selected, community-based 949 [elderly adults](#) with a corrected [serum](#) sodium level between 135.0 and 145.0 mEq/L from the Korean Longitudinal Study on Health and Aging cohort. Patients were stratified by sodium level: 73 in Group 1 (sodium 135.0 to 138.0 mEq/L), 635 in Group 2 (sodium 138.1 to 142.0 mEq/L), and 241 in Group 3 (sodium 142.1 to 145.0 mEq/L).

The researchers found that deaths significantly varied by group: 34 deaths in Group 1, 124 in Group 2, and 52 in Group 3 (P 138.0 mEq/L had a better survival rate than those with a measured sodium level of  $\leq 138.0$  mEq/L and a corrected sodium level of  $\leq 138.0$  mEq/L).

"Sodium level corrected according to serum glucose concentration was a more meaningful risk factor than measured [sodium level](#)," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Low-normal sodium deemed major risk for mortality in elderly (2016, March 25) retrieved 11 February 2023 from <https://medicalxpress.com/news/2016-03-low-normal-sodium-deemed-major-mortality.html>

|   |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|