

# Fatty liver disease is common

April 11 2016

---

Researchers have characterized the prevalence and risk factors of fatty liver disease in patients who undergo liver transplantation. The findings, which are published in *Liver Transplantation*, a journal of the American Association for the Study of Liver Diseases, could have important implications for safeguarding transplant recipients' health.

Steatosis—when the liver becomes infiltrated with fat—occurs frequently after [liver transplantation](#), but little is known about its actual prevalence in transplant recipients, which risk factors are involved, or what effects the condition has on patients' survival. To investigate, Irena Hejlova, MD, of the Institute for Clinical and Experimental Medicine in Prague, and her colleagues retrospectively examined liver biopsies and patient survival data on 548 liver transplant recipients.

Steatosis was found in 309 patients (56.4 percent), including 93 patients (17.0 percent) with significant steatosis. The prevalence increased from 30.3 percent at 1 year to 47.6 percent at 10 years after liver transplantation. When the investigators looked at patients' pre-transplant characteristics, they found that high body mass index and cirrhosis caused by [alcohol consumption](#) were linked with an increased risk of developing steatosis. Regarding patients' post-transplant characteristics, high [body mass index](#), elevated blood triglycerides, alcohol consumption, and type 2 diabetes were linked with steatosis. Post-transplant steatosis was not associated with worse patient survival within several years, but the long-term survival of patients with significant steatosis tended to be worse.

"Our study is the first to describe the occurrence, evolution, and significance of fatty liver in a large unselected population of European liver transplant recipients," said Dr. Hejlova. "Occurrence of fatty liver could indicate those patients whose long-term survival might be impaired. Therefore obesity should be avoided and patients should be encouraged to increase their physical activity. Screening and treatment of diabetes and hyperlipidemia should also be performed, and naturally, liver transplant recipients should abstain from alcohol consumption."

**More information:** Irena Hejlova et al. Prevalence and Risk Factors of Steatosis After Liver Transplantation and Patient Outcomes, *Liver Transplantation* (2015). [DOI: 10.1002/lt.24393](https://doi.org/10.1002/lt.24393)

Provided by Wiley

Citation: Fatty liver disease is common (2016, April 11) retrieved 15 December 2022 from <https://medicalxpress.com/news/2016-04-fatty-liver-disease-common.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.