

# Protein-trapped sugar compounds nourish infant gut microbes

15 April 2016

UC Davis researchers have shown that an enzyme produced by beneficial microbes in babies' intestines is able to harvest specific sugar compounds from human breast-milk and cow's milk. The discovery identifies those sugars—rather than associated protein compounds—as the key to nourishing those important, health-promoting microbes.

"These [sugar](#) compounds selectively provide incredible nutrient support specifically for the growth of the infant gut microbes," said lead researcher David Mills, a professor of food science and technology at UC Davis.

The study results will appear April 15, 2016 online in *Applied and Environmental Microbiology*, a journal of the American Society of Microbiology.

## Advancing earlier discoveries:

In previous studies, Mills and colleagues had shown that [glycoproteins](#) from human milk may be a source of sugar for the [beneficial microbes](#) in the infant gut. Glycoproteins are compounds with both protein and sugar molecules called oligosaccharides.

The researchers also had earlier demonstrated that one such gut microbe—a bacterial subspecies called *Bifidobacterium longum* subsp. *infantis* (*B. infantis*)—produced an enzyme called EndoBI-1 that could split the oligosaccharides away from the glycoproteins.

But there was no definitive answer as to whether it was the sugar or the protein components of the glycoproteins that were supporting growth of *B. infantis*.

## New findings:

In the new study, the researchers set out to determine whether the oligosaccharides could, as

they suspected, be a sole nutrient source for *B. infantis* in the infant gut. They used the EndoBI-1 enzyme to separate the oligosaccharides from the glycoproteins found in bovine colostrum—the "first milk" produced by a lactating cow. Cow's milk was used for the study because it was readily available in significant volumes.

The researchers found that oligosaccharides separated out from the cow's colostrum fueled rapid growth of the *B. infantis* bacterium. However, the oligosaccharides from the cow's milk did not support growth of a related bacterium found in the intestines of adult humans.

The study also showed that *B. infantis* did not grow on milk proteins from which the sugars had been separated out, clearly indicating that the oligosaccharides were the compounds that fed those specific bacteria.

Mills noted that *B. infantis* has many genes that are involved with breaking down glycoproteins in mother's milk in order to release the oligosaccharides.

"Mother's milk co-evolved over millions of years with mammals and the beneficial gut microbiota that the milk helped nourish," he said. "And milk is the only food that co-evolved with humans to make us healthy."

In addition to solving the protein-or-sugar question related to the role of glycoproteins in feeding *B. infantis*, the findings also suggest that cow's [milk](#) could be a source of selective oligosaccharides, which might be used to make therapeutic prebiotics for infants.

Provided by UC Davis

APA citation: Protein-trapped sugar compounds nourish infant gut microbes (2016, April 15) retrieved 27 May 2022 from <https://medicalxpress.com/news/2016-04-protein-trapped-sugar-compounds-nourish-infant.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*