

Blood pressure diet may help prevent gout flares

August 15 2016

New research indicates that a healthy diet can effectively lower blood levels of uric acid, a known trigger of gout. The findings are published in *Arthritis & Rheumatology*, a journal of the American College of Rheumatology (ACR).

Elevated uric acid in the blood plays a key role in gout, an extremely painful form of arthritis that results in profound disability and healthcare expenses. Diet has long been identified as an important determinant of blood uric acid levels, but there is virtually no clinical trial evidence to inform food choice by physicians and patients.

Stephen Juraschek, MD, PhD, of the Johns Hopkins University School of Medicine, and his colleagues looked at the potential of the Dietary Approaches To Stop Hypertension (DASH) diet, a diet with well-established benefits for lowering blood pressure, for lowering uric acid. The DASH diet emphasizes fruits, vegetables, and low-fat dairy foods and reduced consumption of saturated fat, total fat, and cholesterol. It also contains whole grains, lean meats, fish, nuts, and beans.

The investigators assessed a randomized, crossover feeding trial in 103 adults with pre- or stage 1 hypertension. Participants were randomly assigned to receive either the DASH diet or a control diet (typical of the average American diet) and were further fed low, medium, and high sodium levels for 30 days, each in random order.

The DASH diet lowered uric acid on average by 0.35 mg/dL. In

individuals with uric acid levels >7 mg/dL however, which is common among patients with gout, the DASH diet lowered uric acid by >1 mg/dL. While the researchers hypothesized that reducing sodium intake would lower uric acid levels, they found that the opposite was true: higher sodium intake (which was about equal to the average sodium consumed in a typical American diet) decreased uric acid levels compared with low sodium intake. The mechanism by which increased sodium intake decreases uric acid is unclear.

The findings suggest that the DASH diet may represent an effective, non-pharmacologic approach to prevent flares in patients with gout.

"Physicians may now confidently recommend the DASH diet to patients with gout in order to lower uric acid levels," said Dr. Juraschek. "Our findings also show how sodium, or salt, can alter uric acid levels, which provides important insights in further understanding dietary triggers of gout flares."

The research was supported by cooperative agreements and grants from the National Heart, Lung, and Blood Institute and by the General Clinical Research Center Program of the National Center for Research Resources.

More information: Stephen P. Juraschek et al, Effects of the Dietary Approaches To Stop Hypertension (DASH) Diet and Sodium Intake on Serum Uric Acid, *Arthritis & Rheumatology* (2016). DOI: [10.1002/art.39813](https://doi.org/10.1002/art.39813)

Provided by Wiley

Citation: Blood pressure diet may help prevent gout flares (2016, August 15) retrieved 4 May 2023 from <https://medicalxpress.com/news/2016-08-blood-pressure-diet-gout-marker.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.