


Conversion of brain cells offers hope for Parkinson's patients

11 April 2017



Dopamine neurons degenerate and die in the brains of people suffering from Parkinson's disease. Credit: iStock

Researchers at Karolinska Institutet have made significant progress in the search for new treatments for Parkinson's disease. By manipulating the gene expression of non-neuronal cells in the brain, they were able to produce new dopamine neurons. The study, performed on mice and human cells, is published in the prestigious scientific journal *Nature Biotechnology*.

By reprogramming <u>cells</u> in the brain the scientists were able to revert motor symptoms in a mouse model of Parkinson's disease and reproduce the cellular function and responses in human brain cells in the laboratory. The key to this discovery lies in the conversion of brain cells into a subset of nerve cells called <u>dopamine neurons</u>.

Dopamine neurons degenerate and die in the brains of people suffering from Parkinson's disease. The research team showed that they can convert non-neuronal so called <u>glial cells</u> in the brains of mice into new dopamine neurons, in essence, creating new neurons. The glial cells used in the study were astrocytes, named for their star-shape, which are highly abundant in the brains of mice and humans. Using a variety of different transcription factors – proteins that can alter the <u>gene expression</u> – the researchers changed astrocytes in a <u>mouse model</u> of Parkinson's disease into functioning dopamine neurons. This resulted in measurable markers of dopamine signaling in the mouse brain as well as a significant reduction in the symptoms of the disease.

The team also performed experiments on human astrocytes in the laboratory which they were able to convert into functional dopamine neurons, raising the possibility for treatment of patients suffering from Parkinson's disease.

An alternative to cell therapy

Cell replacement therapies for neurodegenerative disease have focused on transplantation of the cell types affected by the pathological process.

"We think that in the future, it may be possible to deliver genes and small molecules, rather than cells, to replace the dopamine neurons lost in Parkinson's disease" says Professor Ernest Arenas at Karolinska Institutet's Department of medical biochemistry and biophysics.

More information: Pia Rivetti di Val Cervo et al. Induction of functional dopamine neurons from human astrocytes in vitro and mouse astrocytes in a Parkinson's disease model, *Nature Biotechnology* (2017). DOI: 10.1038/nbt.3835

Provided by Karolinska Institutet



APA citation: Conversion of brain cells offers hope for Parkinson's patients (2017, April 11) retrieved 18 April 2021 from <u>https://medicalxpress.com/news/2017-04-conversion-brain-cells-parkinson-patients.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.