

IgE allergy testing improves atopic dermatitis outcomes

July 6 2017



(HealthDay)—Identification of allergens by immunoglobulin E (IgE)



testing improves outcomes in atopic dermatitis, according to a study published online June 20 in the *International Journal of Dermatology*.

Brett M. Will, from Georgetown University in Washington, D.C., and colleagues retrospectively assessed affected body surface area at first presentation, IgE quantitative assaying of allergens (IgEQAA) classes, and total IgE concentration for 54 patients with atopic dermatitis.

The researchers found that 41 of 54 patients had an abnormally high total IgE concentration (76 percent). After making <u>lifestyle changes</u>, nine (17 percent) of these patients showed significant improvement in their symptoms.

"While IgEQAA has historically been used by allergists, it has yet to see the same widespread usage within the field of dermatology. IgEQAA is easy to perform compared to skin prick testing and may be helpful in some patients who are not responding to conservative treatment," the authors write. "Further studies are required; however, from our data it appears that IgEQAA testing can have a significant impact on treatment outcomes in selected patients with <u>atopic dermatitis</u>."

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

Citation: IgE allergy testing improves atopic dermatitis outcomes (2017, July 6) retrieved 19 November 2023 from <u>https://medicalxpress.com/news/2017-07-ige-allergy-atopic-dermatitis-outcomes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.