


Ozone pollution connected to cardiovascular health

17 July 2017



Human heart. Credit: copyright American Heart Association

Exposure to ozone, long associated with impaired lung function, is also connected to health changes that can cause cardiovascular disease such as heart attack, high blood pressure and stroke, according to a new study of Chinese adults.

These findings, by a team from Duke University, Tsinghua University, Duke Kunshan University and Peking University, appear in the July 17, 2017 edition of *JAMA Internal Medicine*.

Ozone is a pollutant formed through a chemical reaction that occurs when sunlight interacts with nitrogen oxides and other organic compounds that are generated by coal-burning, vehicle exhaust and some natural sources.

"We know that [ozone](#) can damage the respiratory

system, reduce [lung function](#) and cause asthma attacks," said study author Junfeng (Jim) Zhang, from Duke and Duke Kunshan University. "Here, we wanted to learn whether ozone affects other aspects of human health, specifically the cardiovascular system."

Zhang and colleagues studied 89 healthy adults living in Changsha City, China, for one year. They monitored indoor and outdoor [ozone levels](#), along with other pollutants. At four intervals, the study team took participant blood and urine samples and used a breathing test called spirometry to examine a set of factors that could contribute to cardiovascular and respiratory disease.

The team examined inflammation and oxidative stress, arterial stiffness, [blood pressure](#), clotting factors and lung function in participants. They noted blood platelet activation (a risk factor for clotting) and an increase in blood pressure, suggesting a possible mechanism by which ozone may affect cardiovascular health. These effects were found with ozone exposure lower than that which affects respiratory health, and lower than current Environmental Protection Agency (EPA) air quality standards.

"This study shows that standards for safe ozone exposure should take into account its effect on [cardiovascular disease](#) risk," said Zhang.

"In 2015, 108 million Americans—one third of the population—lived in counties with ozone levels that exceeded standards set by the EPA," Zhang said. "In contrast, only 31 million Americans live in counties where other pollutants exceed EPA standards."

The production of ozone globally will be exacerbated by a warmer climate, "so it will be an increasing trend with climate change," said Zhang. Ozone is a difficult pollutant to control because its creation in the atmosphere is complex. "For

example, a reduction in [nitrogen oxides](#) does not necessarily mean a reduction in ozone levels," Zhang said.

More information: "Pathophysiologic Mechanisms Underlying Cardiorespiratory Effects of Ozone in healthy Adults," Drew B Day, Jianbang Xiang, Jinhan Mo, Feng Li, Mingkei Chung, Jicheng Gong, Charles Weschler, Pamela A Ohman-Strickland, Jan Sundell, Wenguo Weng, Yinping Zhang Junfeng (Jim) Zhang. *JAMA Internal Medicine*, July 17, 2017.

Provided by Duke University

APA citation: Ozone pollution connected to cardiovascular health (2017, July 17) retrieved 13 November 2022 from <https://medicalxpress.com/news/2017-07-ozone-pollution-cardiovascular-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.