


Zika virus infects developing brain by first infecting cells meant to defend against it

October 27 2017



A false color micrograph shows microglia cells (green) infected by the Zika virus (blue). Credit: UC San Diego Health

Researchers at the University of California San Diego School of Medicine, with colleagues in Brazil, report that the Zika virus is transmitted from mother to fetus by infected cells that, ironically, will later develop into the brain's first and primary form of defense against invasive pathogens.

The findings are published in the current online issue of *Human Molecular Genetics*.

"It's a Trojan Horse strategy," said Alysson Muotri, PhD, professor in the UC San Diego School of Medicine departments of Pediatrics and Cellular and Molecular Medicine. "During embryogenesis—the early stages of prenatal development—[cells](#) called microglia form in the yolk sac and then disperse throughout the central nervous system (CNS) of the developing child.

"In the brain, these microglia will become resident macrophages whose job is to constantly clear away plaques, damaged cells and infectious agents. Our findings show that the Zika virus can infect these early microglia, sneaking into the brain where they transmit the virus to other brain cells, resulting in the devastating neurological damage we see in some newborns."

Beginning in 2015, a dramatic increase in children born with microcephaly or abnormally small heads and other birth defects was observed in Brazil. The phenomenon was subsequently linked to infection by the Zika virus, which Muotri and others confirmed last year caused birth defects in experimental models.

Typically, the Zika virus is transmitted to people through the bite of infected *Aedes* species mosquitoes. However, a pregnant woman can also pass the virus to her fetus, though scientists have not been able to precisely describe the mode of transmission.

"Considering the timing of transmission, we hypothesized that microglia might be serving as a Trojan horse to transport the virus during invasion of the CNS," Muotri said.

To test their hypothesis, the researchers used human induced [pluripotent](#)

[stem cells](#) to create two relevant CNS [cell types](#): microglia and [neural progenitor cells](#) (NPCs), which generate the millions of neurons and glial cells required during embryonic development. Then they established a co-culture system that mimicked the interactions of the two cell types *in vitro* when exposed to the Zika virus.

They found that the microglia cells engulfed Zika-infected NPCs, doing their job. But when these microglia carrying the virus were placed in contact with non-infected NPCs, they transmitted the virus to the latter. "That suggests microglia may indeed be the culprit for transmitting the virus to the CNS during prenatal neurodevelopment," Muotri said.

Muotri and colleagues then tested whether an FDA-approved drug called Sofosbuvir, marketed as Sovaldi and used to treat hepatitis C, might limit viral infection of NPCs in co-culture with infected [microglia](#). It did. "Sofosbuvir significantly decreased cell death of NPCs and the viral load in NPCs"

Though the findings are based on *in vitro* research and further investigation is necessary, Muotri said they were encouraging, suggesting microglial cells could be a therapeutic target for reducing Zika transmission into the CNS of developing fetuses.

"The co-culture system we've developed is robust and useful for studying neuro-immune interactions," he said. "It can also serve as a drug-screening platform for discovering new therapeutic compounds against Zika [virus](#) infections in a human context."

More information: Pinar Mesci et al, Modeling neuro-immune interactions during Zika virus infection, *Human Molecular Genetics* (2017). [DOI: 10.1093/hmg/ddx382](https://doi.org/10.1093/hmg/ddx382)

Provided by University of California - San Diego

Citation: Zika virus infects developing brain by first infecting cells meant to defend against it (2017, October 27) retrieved 4 February 2024 from

<https://medicalxpress.com/news/2017-10-zika-virus-infects-brain-infecting.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.