


Higher estrogen levels linked to increased alcohol sensitivity in brain's 'reward center'

November 6 2017



Amy Lasek, assistant professor of psychiatry, UIC College of Medicine. Credit: Jenny Fontaine.

The reward center of the brain is much more attuned to the pleasurable effects of alcohol when estrogen levels are elevated, an effect that may

underlie the development of addiction in women, according to a study on mice at the University of Illinois at Chicago.

Led by Amy Lasek, assistant professor of psychiatry in the UIC College of Medicine, researchers found that [neurons](#) in a region of the brain called the ventral tegmental area, or VTA (also known as the "[reward center](#)"), fired most rapidly in response to alcohol when their estrogen levels were high. This response, according to their findings published online in the journal *PLOS ONE*, is mediated through receptors on dopamine-emitting neurons in the VTA.

"When estrogen levels are higher, alcohol is much more rewarding," said Lasek, who is the corresponding author on the paper and a researcher in the UIC Center for Alcohol Research in Epigenetics. "Women may be more vulnerable to the effects of alcohol or more likely to overindulge during certain stages of their cycle when estrogen levels are higher, or may be more likely to seek out alcohol during those stages."

Studies indicate that gender differences in psychiatric disorders, including addiction, are influenced by estrogen, one of the primary female sex hormones. Women are more likely to exhibit greater escalation of abuse of alcohol and other drugs, and are more prone to relapse in response to stress and anxiety.

The VTA helps evaluate whether something is valuable or good. When neurons in this area of the brain are stimulated, they release dopamine—a powerful neurotransmitter responsible for feelings of wellness—and, in large doses, euphoria. When something good is encountered—for example, chocolate—the neurons in the VTA fire more rapidly, enforcing reward circuitry that encodes the idea that chocolate is enjoyable and something to be sought out. Over time, the VTA neurons fire more quickly at the sight, or even thought of, chocolate, explained Lasek. In addiction, VTA neurons are tuned into

drugs of abuse, and fire more quickly in relation to consuming or even thinking about drugs, driving the person to seek them out—often at the expense of their own health, family, friends and jobs.

Many animal studies have shown that alcohol increases the firing of dopamine-sensitive neurons in the VTA, but little is known about exactly why this occurs.

Lasek and her colleagues examined the relationship between estrogen, alcohol and the VTA in female mice. They used naturally cycling mice that were allowed to go through their normal estrous cycles, akin to the menstrual cycle in women.

Mice were evaluated to determine when they entered diestrus—the phase in the estrous cycle when estrogen levels are close to their peak.

"In mice in diestrus, estrogen levels increase to about 10 times higher than they are in estrus, the phase in which ovulation occurs and estrogen levels drop," Lasek said.

VTAs were taken from mice in both estrus and diestrus and kept alive in special chambers. Electrodes recorded the activity of individual dopamine-sensitive neurons in the VTA. Next, the researchers added alcohol to the chamber. Activity increased twice as much in neurons from mice in diestrus compared to the response of neurons from mice in estrus.

Lasek and her colleagues then blocked estrogen receptors on dopamine-sensitive neurons in VTA in mice in estrus and diestrus. With the blocker present, the response to alcohol in neurons from mice in diestrus was significantly lower compared with neurons where estrogen receptors remained functional. The [estrogen receptor](#) blocker reduced the alcohol response to levels seen in mice in estrus. The responses to alcohol in

neurons from [mice](#) in estrus were unaffected by the estrogen receptor blocker.

"The increased reward response to alcohol we see when estrogen levels are high is mediated through receptors for estrogen in the VTA," said Mark Brodie, professor of physiology and biophysics in the UIC College of Medicine and a co-author on the paper.

Lasek believes that the increased sensitivity to alcohol in the VTA when [estrogen levels](#) peak may play a significant role in the development of addiction in women.

"We already know that binge drinking can lead to lasting changes in the brain, and in women, those changes may be faster and more significant due to the interaction we see between alcohol, the VTA and [estrogen](#)," Lasek said. "Binge drinking can increase the risk of developing alcoholism, so women need to be careful about how much alcohol they drink. They should be aware that they may sometimes inadvertently over-consume alcohol because the area of the brain involved in [alcohol](#) reward is responding very strongly."

More information: Bertha J. Vandegrift et al, Estradiol increases the sensitivity of ventral tegmental area dopamine neurons to dopamine and ethanol, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0187698](https://doi.org/10.1371/journal.pone.0187698)

Provided by University of Illinois at Chicago

Citation: Higher estrogen levels linked to increased alcohol sensitivity in brain's 'reward center' (2017, November 6) retrieved 13 February 2024 from <https://medicalxpress.com/news/2017-11-higher-estrogen-linked-alcohol-sensitivity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.