

Cannabis linked to bipolar symptoms in young adults

30 November 2017



Credit: CC0 Public Domain

Cannabis use in youth is linked to bipolar symptoms in young adults, finds new research by the University of Warwick.

Researchers from Warwick Medical School found that adolescent [cannabis](#) use is an independent risk factor for future hypomania - periods of elated mood, over-active and excited behaviour, and reduced need for sleep that are often experienced as part of [bipolar disorder](#), and have a significant impact on day-to-day life.

Led by Dr Steven Marwaha, a clinical academic Psychiatrist, the research analysed data from the Avon Longitudinal Study of Parents and Children and found that teenage cannabis use at least 2-3 times weekly is directly associated with suffering from symptoms of hypomania in later years.

There was a dose response relationship such that any use still increased the risk but less powerfully.

The Warwick research is the first to test the prospective association between adolescent cannabis use and hypomania in early adulthood,

whilst controlling for important other factors that might explain this connection (e.g psychotic symptoms).

Cannabis use was also found to mediate the association of both childhood sexual abuse and hypomania, and male gender and hypomania.

The findings suggest frequent [adolescent](#) cannabis use is likely to be a suitable target for interventions that may allay the risk of young people developing bipolar disorder.

Commenting on the research, Dr Marwaha said: "Cannabis use in [young people](#) is common and associated with psychiatric [disorders](#). However, the prospective link between cannabis use and bipolar disorder symptoms has rarely been investigated.

"Adolescent cannabis use may be an [independent risk factor](#) for future hypomania, and the nature of the association suggests a potential causal link. As such it might be a useful target for indicated prevention of hypomania."

Cannabis is one of the most commonly used illegal substances of abuse in western countries. Problematic use in the general population is as high as 9.5% in the United States, while 2.6% of the UK population report having been cannabis dependent in the last year.

More information: Cannabis Use and Hypomania in Young People: A Prospective Analysis, *Schizophrenia Bulletin* (2017). [DOI: 10.1093/schbul/sbx158](#)

Provided by University of Warwick

APA citation: Cannabis linked to bipolar symptoms in young adults (2017, November 30) retrieved 10 September 2022 from <https://medicalxpress.com/news/2017-11-cannabis-linked-bipolar-symptoms-young.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.