

Significant ultrasound practice needed to diagnose appendicitis

18 January 2018



90.9 and 93.5 percent, respectively, after the first 20 cases.

"A minimum of 20 cases are needed to achieve competency in [emergency](#) ultrasonography diagnosis of [acute appendicitis](#)," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—There is a learning curve among novice emergency medicine residents in using emergency ultrasonography to diagnose acute appendicitis, according to a study published online Jan. 5 in the *Journal of Clinical Ultrasound*.

Jeehyun Kim, M.D., from the Chungnam National University Hospital in Daejeon, South Korea, and colleagues compared agreement between the appendicitis diagnoses of three novice emergency medicine residents who completed a one-day training course and an experienced, board-certified radiologist. (Some patients also underwent computed tomography.)

The researchers note that a total of 266 patients were included, and the overall Cohen's kappa coefficient was 0.77. For the first 20 cases, the kappa value was 0.49, but it increased rapidly during this time, and the kappa coefficient was 0.84 after the first 20 cases. The sensitivity and specificity values were 64.3 and 84.4 percent, respectively, for the first 20 cases, but increased to

APA citation: Significant ultrasound practice needed to diagnose appendicitis (2018, January 18) retrieved 10 November 2022 from <https://medicalxpress.com/news/2018-01-significant-ultrasound-appendicitis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.