

# Healthy diet may not offset high salt intake

5 March 2018



Credit: CC0 Public Domain

A healthy diet may not offset the effects of a high salt intake on blood pressure, suggests a new study.

The research, from scientists at a number of institutions, including Imperial College London and Northwestern University, analysed the diets of over 4,000 people.

The results, published in the journal *Hypertension*, showed that people eating higher amounts of salt had higher blood pressure - no matter how healthy a person's overall diet.

The scientists behind the research are now advising people to monitor their [salt intake](#) - and for food manufacturers to lower the [salt content](#) in their products.

High blood pressure affects more than one in four adults in the UK, and increases the risk of a number of conditions including heart attacks and stroke. It's thought to have a number of causes, including age, weight and eating too much salt.

It's thought that vitamins and minerals in fruit and vegetables might in some way affect blood

vessels, enabling them to [lower blood pressure](#).

Previously, experts believed that eating high amounts of fruit and vegetables might help counteract the effect of high salt on blood pressure.

However while these foods do tend to lower blood pressure, the new research suggests they do not counteract the adverse influence of salt intake.

In the paper, the team studied data from the so-called INTERMAP study. In this study, which was conducted between 1997-1999, scientists tracked the diets of 4,680 people, aged 40-59, from the USA, UK, Japan and China. The volunteers were tracked over four days, and two urine samples were taken during this time. Measurements of height, weight and blood pressure were also taken. The study data has since been used for numerous research projects.

In the latest paper, published today, the team assessed concentrations of sodium and potassium in the [urine samples](#). Sodium is the main component of salt, while potassium, which is found in green leafy vegetables, has been linked to lower blood pressure.

The team also used dietary data to assess the volunteers' intake of over 80 nutrients that may be linked to low blood pressure, including vitamin C, fibre, and omega-3 fatty acids. Many of these nutrients are found in fruit, vegetables and whole grains.

The researchers found a correlation between high blood pressure and higher salt intake, even in people who were eating a high amount of potassium and other nutrients. The researchers estimated salt intake by analysing sodium in the urine, as well as analysing dietary data.

The recommended upper limit of adult salt intake in the UK is 6g a day - around one teaspoon.

The study found that average salt intake across the

study was 10.7g a day. The average intake for the UK was 8.5g, while the intake for the USA, China and Japan were 9.6g, 13.4g and 11.7g respectively.

Provided by Imperial College London

Increasing salt intake above this average amount was linked to an increased in blood pressure. An increase of an additional 7g (1.2 teaspoons) of salt above the average intake was associated with an increase in [systolic blood pressure](#) of 3.7 mmHg.

Blood pressure is measured in two numbers - the first, called systolic pressure, measures the force the heart pumps blood around the body. The second number, called diastolic pressure, is the resistance to blood flow in the arteries. Ideally, blood pressure should be between 90/60 and 120/80 mmHg. However, reducing blood pressure by just a small amount can reduce the risk of conditions such as stroke.

Dr Queenie Chan, joint lead author of the research from the School of Public Health at Imperial, said the research shows the importance of cutting salt intake.

"We currently have a global epidemic of high salt intake - and [high blood pressure](#). This research shows there are no cheats when it comes to reducing [blood pressure](#). Having a low salt diet is key - even if your diet is otherwise healthy and balanced."

She added: "As a large amount of the salt in our diet comes from processed food, we are urging food manufacturers to take steps to reduce [salt](#) in their products."

The team acknowledge that because the data was collected over four days, it provides information from a snapshot of time. They now hope to focus on longer term studies, with a greater number of people.

**More information:** *Hypertension* (2018).  
[dx.doi.org/10.1161/HYPERTENSIONAHA.117.09928](https://doi.org/10.1161/HYPERTENSIONAHA.117.09928)

APA citation: Healthy diet may not offset high salt intake (2018, March 5) retrieved 19 September 2022 from <https://medicalxpress.com/news/2018-03-healthy-diet-offset-high-salt.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*