

Food insecurity tied to poor glycemic control in diabetes

23 March 2018



"Food insecurity is associated with higher HbA1c, but living in an area with low physical food access is not," the authors write. "Food insecurity screening and interventions may help improve glycemic control for vulnerable patients."

More information: [Abstract/Full Text](#)
(subscription or payment may be required)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Limited food access owing to cost (food insecurity) is associated with increased hemoglobin A1c (HbA1C) among patients with diabetes, according to a study published online March 19 in *Diabetes Care*.

Seth A. Berkowitz, M.D., from Massachusetts General Hospital in Boston, and colleagues examined the correlation between [food insecurity](#), low physical food access, and [glycemic control](#) in a random sample of patients with diabetes in a primary care network. A total of 391 participants were followed for a mean of 37 months.

The researchers found that 20 and 31 percent of respondents reported food insecurity and lived in an area of low physical food access, respectively. Food insecurity was correlated with increased HbA1c (difference, 0.6 percent) in adjusted models, which did not improve over time. Living in an area with low physical food access was not linked to a difference in HbA1c or with change over time.

APA citation: Food insecurity tied to poor glycemic control in diabetes (2018, March 23) retrieved 10 June 2021 from <https://medicalxpress.com/news/2018-03-food-insecurity-tied-poor-glycemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.