


Revolutionary brain-mapping technique provides new blueprint for cortical connections

March 28 2018




The large, triangular-shaped central area identifies the primary visual cortex in mouse. By barcoding 591 neurons in this region, the team used MAPseq to discover patterns of their projections to nearby higher visual cortical areas. Each of the latter is identified in this calibration image, which registers the brain's reaction to two different kinds of visual stimuli (registering in pink and green fluorescence). Credit: Zador Lab, CSHL

Using a revolutionary new brain-mapping technology recently developed at Cold Spring Harbor Laboratory (CSHL), an international team of scientists led by Professor Anthony Zador have made a discovery that will force neuroscientists to rethink how areas of the cortex communicate with one another.

The [new technology](#), called MAPseq, allowed the scientists to determine that neurons in the [primary visual cortex](#) communicate with higher visual areas of the cortex much more broadly than previously believed, and according to specific patterns.

The wiring diagram of the cortex determines how information is processed across dozens of [cortical areas](#). "If we don't know how information is combined even at the earliest stages, then we have essentially no shot of figuring out how the brain works." says Justus Kebschull, now a postdoctoral researcher at Stanford University who was instrumental in developing MAPseq as a graduate student in Zador's lab.

In a MAPseq experiment, hundreds or thousands of neurons are labeled uniquely with random RNA sequences ("barcodes") via a single injection made in the brain. The barcodes are transported into the branching axons of each labeled neuron, where they can later be read out by high-throughput barcode sequencing after the brain is dissected. This process allows researchers to identify every brain area that each barcoded neuron makes contact with.



Left: Single-neuron tracing reveals the projections of a single neuron in the mouse brain. It took the team 3 years to map 31 such neurons. In 3 weeks and at a fraction of the cost, MAPseq discovered the destinations of 591 neurons in the mouse primary visual cortex, reaching similar conclusions about projection patterns. The 'readout' is dramatically different (right), showing 8 projection 'motifs' (8 clusters, read horizontally), with strength of connections in parts of the higher visual cortex displayed in color intensities. Credit: Zador Lab, CSHL

The Zador lab has begun to use MAPseq to compare the brains of a mouse autism model with healthy mouse brains to see if mis-wiring occurs during development at the single-neuron level that might explain the disorder's symptoms—just one of many potential applications of the method.

In the experiments reported today, the team first verified the relatively new technology by comparing its results with the gold-standard mapping method, called single-neuron tracing. They used the latter method to trace 31 mouse neurons in the primary visual cortex to up to seven different cortical locations. These experiments took 3 years to complete. It then took them only 3 weeks to use MAPseq to map the projections within the cortex arising from 591 primary visual cortex neurons. Both techniques revealed that the majority of neurons in this location target multiple visual areas, and that roughly the same proportions of cells project to two, three, or four cortical areas.

Beyond just being faster, MAPseq was able to reveal patterns in the way that [neurons](#) in the primary visual cortex connected to other [visual areas](#) across the [cortex](#). About 73% fit into one of six distinct projection patterns. This architecture may facilitate coordination of activity among those areas, which in turn could provide a means of linking visual information across the brain to help form complex percepts.

"Our finding signals a shift away from the rather convenient idea of every neuron projecting to just one cortical area," says Kebschull. "That thinking ignores the underlying structure of the [brain](#), and in the future, the way people do their experiments is going to change drastically."

Han Y et al, "A single-cell anatomical blueprint for intracortical communication" appears online March 28, 2018 in *Nature*.

More information: "A single-cell anatomical blueprint for intracortical communication" *Nature* (2018).

Provided by Cold Spring Harbor Laboratory

Citation: Revolutionary brain-mapping technique provides new blueprint for cortical connections (2018, March 28) retrieved 2 April 2023 from <https://medicalxpress.com/news/2018-03-revolutionary-brain-mapping-technique-blueprint-cortical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.