


Study: Almost 100 million adults have COPD in China

10 April 2018



Micrograph showing emphysema (left – large empty spaces) and lung tissue with relative preservation of the alveoli (right). Credit: Wikipedia, CC-BY-SA 3.0

Chronic obstructive pulmonary disease (COPD) is widespread in China with 8.6 percent of the country's adult population - almost 100 million people -suffering from the chronic lung disease, according to a new Tulane University study published in *The Lancet*.

The study, which provided lung-function screenings for more than 50,990 participants, is the largest survey of COPD across age groups ever conducted in China, researcher say.

COPD, an inflammatory lung disease that causes obstructed airflow into the lungs, is the third leading cause of death in China. It is caused by long-term exposure to irritants in the air, including [cigarette smoke](#). During the past decade, ambient air pollution has become a major public-health crisis in the country while cigarette smoking remains high, especially among men, says senior author Dr. Jiang He, Joseph S. Copes Chair of Epidemiology at Tulane University School of Public Health & Tropical Medicine.

"Our research shows that COPD is highly prevalent in the Chinese [adult population](#). The biggest preventable risk factors for the disease are cigarette smoking and air pollution," He says. "Prevention programs and increased efforts to catch COPD early should be public-health priorities in China to reduce COPD-related diseases and deaths."

The study found that men had a higher prevalence (11.9 percent) of COPD than women (5.4 percent). The sex difference was noted among all age groups in the general population and even among those who never smoked.

Based on the lung-function screenings, researchers estimated the total number of individuals aged 20 years or older with COPD in China was 99.9 million in 2015. Of these, 68.4 million were men and 31.5 million were women.

Researchers found that most people with COPD were unaware of their condition, and few had received a pulmonary function test prior to the study. Only 2.6 percent of participants were aware of their condition, and only 12 percent of those found to have COPD reported getting a previous [pulmonary function test](#).

Researchers found COPD prevalence increased with age. It was 2.1 percent for individuals aged 20-39 years old compared to 13.7 percent for those aged 40 years or older. The [disease](#) was more common for rural residents (9.6 percent) than for those in urban areas (7.4 percent). The same trend was noted among non-smokers.

Researchers suspect that those who didn't smoke developed COPD due to other risk factors or environmental exposures such as air pollution, chronic cough during childhood, parental history of respiratory diseases, low bodyweight and low education.

The study calls for new national public health strategies to prevent COPD that include smoking cessation programs, stricter control of ambient [air pollution](#) and biomass use, and COPD screenings for high-risk individuals using pulmonary function tests.

Provided by Tulane University

APA citation: Study: Almost 100 million adults have COPD in China (2018, April 10) retrieved 23 June 2022 from <https://medicalxpress.com/news/2018-04-million-adults-copd-china.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.