

Screening for impaired vision in older adults: New Canadian guideline

May 14 2018

A new Canadian guideline for impaired vision in older adults recommends against primary care screening of older adults not reporting concerns about their vision. The guideline, published in English and French in *CMAJ* (*Canadian Medical Association Journal*) by the Canadian Task Force on Preventive Health Care (CTFPHC), is aimed at primary care practitioners.

Visual impairment describes less than 20/40 vision, which usually cannot be corrected with glasses, contact lenses or vision-related procedures. This level of difficulty with vision can affect quality of life, as well as participation in work, social and leisure activities, and increases the likelihood of injuries from falls and other accidents.

The <u>task force</u> considered the benefits of screening for visual impairment in primary health care and referring patients to optometrists for formal vision testing.

"We found no evidence of benefit to patients aged 65 years or older from being screened for impaired vision as a way to prevent limitations on daily living or other consequences," said Dr. Brenda Wilson, Task Force Impaired Vision Working Group Chair. "The task force therefore recommends against screening for impaired vision in primary care settings for people living independently in the community."

Currently, people must make their own appointments for regular vision screening or if they suspect visual problems. Most provinces in Canada



cover comprehensive eye examinations for adults aged 65 years and older by eye care professionals.

The new guideline updates a previous guideline from 1995, which recommended screening for visual impairment in elderly patients with diabetes of at least 5 years' duration. The 2018 guideline is based on the latest and highest-quality evidence on screening, which includes 15 randomized controlled trials involving participants aged 65 years or older. It is consistent with the recommendation on <u>vision</u> screening for <u>older adults</u> from the United States Preventive Services Task Force.

"Although <u>screening</u> does not appear to be effective, we need to look for ways to effectively support older Canadians who do experience <u>visual</u> <u>impairment</u> so that they get the services they need from optometrists or other eye care professionals" said Dr. Brett Thombs, chair of the CTFPHC.

More information: Screening for impaired vision in communitydwelling adults aged 65 years and older in primary care settings, <u>https://doi.org/10.1503/cmaj.171430</u>

Provided by Canadian Medical Association Journal

Citation: Screening for impaired vision in older adults: New Canadian guideline (2018, May 14) retrieved 19 November 2023 from <u>https://medicalxpress.com/news/2018-05-screening-impaired-vision-older-adults.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.