

Newly developed therapeutic shown to combat drug addiction

June 29 2018

A new therapeutic may help reverse chemical imbalances made to the brain by habitual drug use and could one day help recovering drug addicts avoid future drug use.

Researchers from The University of Texas Medical Branch at Galveston developed and tested a treatment on rats and found it effective in reducing the animals' cravings. Their findings are in the *Journal of Medicinal Chemistry*.

When someone habitually misuses drugs, their brain chemistry is changed in ways that make it harder for them to quit taking drugs despite negative consequences. Once someone has developed this brain disorder, their mind pays sharper attention to cues that encourage <u>drug</u> use, making it harder for them to abstain.

Serotonin, a brain chemical that transmits information between neural regions, is a key player in these changes. There are currently no medications available to correct this chemical imbalance.

UTMB pharmacology and toxicology professors Jia Zhou, Kathryn Cunningham and their colleagues found that the serotonin 2C receptors in drug addicts do not work as well as they should. They designed, synthesized and pharmacologically evaluated a series of small molecule therapeutics designed to restore the weakened signaling.

The researchers trained rats to press on a lever for cocaine infusions at



certain light cues. Once the rats learned this cocaine-seeking behavior, half of them received the most promising therapeutic and the other half received only saline.

The animals treated with the new therapeutic pressed the lever for cocaine far fewer times than the saline-treated control animals, even when reinforced with the cocaine-associated light cues.

"We are the first to show that a serotonin 2C receptor therapeutic of this type can be successfully used to decrease drug-seeking behaviors," said Cunningham, the director of UTMB's Center for Addiction Research. "Our findings are especially exciting because in addition to someday helping people to recover from <u>drug addiction</u>, impaired functioning of the <u>serotonin</u> 2C receptor is also thought to contribute to other chronic <u>health issues</u> such as depression, impulsivity disorders, obesity and schizophrenia."

"We will continue to optimize the chemical structure and effectiveness of our new therapeutic and conduct safety testing in cellular and animal studies," said Zhou. "We look forward to the future when we can begin clinical trials so that this therapeutic can begin to help people free themselves from drug addiction and other health issues."

More information: Christopher T. Wild et al. Design, Synthesis, and Characterization of 4-Undecylpiperidine-2-carboxamides as Positive Allosteric Modulators of the Serotonin (5-HT) 5-HT2C Receptor, *Journal of Medicinal Chemistry* (2018). DOI: 10.1021/acs.jmedchem.8b00401

Provided by University of Texas Medical Branch at Galveston



Citation: Newly developed therapeutic shown to combat drug addiction (2018, June 29) retrieved 11 February 2023 from

https://medicalxpress.com/news/2018-06-newly-therapeutic-shown-combat-drug.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.